

Active Recombinant Human TNFRSF10B

Cat. No. TNFRSF10B-518H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human TNFRSF10B produced in E. coli is approximately 14.8 kDa, a single non-glycosylated polypeptide chain containing 132 amino acids.
Species	Human
Source	E.coli
Description	Tumor necrosis factor-related apoptosis-inducing ligand Receptor 2 (TRAIL-R2) is a cell-surface receptor involved in tumor necrosis factor-related apoptosis-inducing ligand (TRAIL)-induced cell-death signaling. 1 The death ligand TRAIL bears high potential as a new anticancer agent, as binding to the death receptors TRAIL-R1 or TRAIL-R2 triggers apoptosis in most cancer cells. 2 TRAIL-R2 has been shown to be associated with a decrease in the survival rates of breast cancer patients.
Form	Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 determined by its ability to inhibit TRAIL-mediated cytotoxicity using mouse L929 cells treated with TRAILI is less than 2 ng/ml, corresponding to a specific activity of $>5.0 \times 10^5$ IU/mg.
Molecular Mass	14.8 kDa
AA Sequence	ESALITQQDL APQQRAPQQ KRSSPSEGLC PPGHHISEDG RDCISCKYGQ DYSTHWNDLL FCLRCTRCDS GEVELSPCTT TRNTVCQCEE GTFREEDSPE MCRKCRTGCP RGMVKVGDC T PWSDIECVHK ES

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Less than 1EU/μg of rHusTRAIL-R2/TNFRSF10B as determined by LAL method.
Purity	>97% by SDS-PAGE and HPLC analyses.
Usage	This material is for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE.
Storage	This lyophilized preparation is stable at 2-8 centigrade, but should be kept at -20 centigrade for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 centigrade. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 centigrade to -70 centigrade. Avoid repeated freeze/thaw cycles.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	TNFRSF10B tumor necrosis factor receptor superfamily, member 10b [Homo sapiens]
Official Symbol	TNFRSF10B
Synonyms	TNFRSF10B; tumor necrosis factor receptor superfamily, member 10b; tumor necrosis factor receptor superfamily member 10B; CD262; DR5; KILLER; TRAIL R2; TRICK2A; TRICKB; Fas-like protein; death receptor 5; cytotoxic TRAIL receptor-2; apoptosis inducing receptor TRAIL-R2; apoptosis inducing protein TRICK2A/2B; TNF-related apoptosis-inducing ligand receptor 2; death domain containing receptor for

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	TRAIL/Apo-2L; tumor necrosis factor receptor-like protein ZTNFR9; p53-regulated DNA damage-inducible cell death receptor(killer); TRICK2; ZTNFR9; TRAILR2; TRICK2B; TRAIL-R2; KILLER/DR5;
Gene ID	8795
mRNA Refseq	NM_003842
Protein Refseq	NP_003833
MIM	603612
UniProt ID	O14763
Chromosome Location	8p22-p21
Pathway	Activation of Pro-Caspase 8, organism-specific biosystem; Apoptosis, organism-specific biosystem; Apoptosis, organism-specific biosystem; Apoptosis, conserved biosystem; Apoptosis, organism-specific biosystem; Caspase-8 is formed from procaspase-8, organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem;
Function	TRAIL binding; protein binding; receptor activity;