

Recombinant Rhesus IL4 protein

Cat. No. IL4-540R **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Rhesus IL4 protein was expressed in Escherichia coli.
Species	Rhesus macaque
Source	E.coli
ProteinLength	129
Description	Interleukin-4 (IL-4) is a pleiotropic cytokine mainly produced by mast cells, activated T cells and bone marrow stromal cells. It has many biological roles, including the stimulation of activated B-cell and T-cell proliferation, and the differentiation of CD4+ T-cells into Th2 cells. Additionally, It enhances both secretion and cell surface expression of IgE and IgG1 and also regulates the expression of the low affinity Fc receptor for IgE (CD23) on both lymphocytes and monocytes. The rhesus macaque IL-4 shares about 40 % ~ 45 % a.a. sequence identity with human, mouse, rat IL-4 and they are species-specific in their activities.
Form	Lyophilized from a 0.2µm filtered concentrated solution in 2 × PBS, pH 7.4, 5 % trehalose.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by a cell proliferation assay using human TF-1 cells is less than 1.0 ng/ml, corresponding to a specific activity of > 1.0 × 10 ⁶ IU/mg.
Molecular Mass	Approximately 14.9 kDa, a single non-glycosylated polypeptide chain containing 129

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	amino acids.
AA Sequence	HNCHIALREIIETLNSLTEQKTLCTKLTITDILAASKNTEKETFCRAATVLRQFYSHHE KDTRCLGATAQQFHRHKQLIRFLKRLDRNLWGLAGLNSCPVKEANQSTLEDFLERL KTIMREKYSKCSS
Endotoxin	Less than 1 EU/μg of rRhIL-4 as determined by LAL method.
Purity	>96% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	IL4
Official Symbol	IL4
Synonyms	BSF-1, Lymphocyte Stimulatory Factor 1
Gene ID	574281

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq [NM_001032904.1](#)

Protein Refseq [NP_001028076.1](#)

UniProt ID [P51492](#)

SDS-PAGE of IL4-
540R



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA