

Recombinant Mouse Mdk protein

Cat. No. Mdk-567M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse Mdk protein was expressed in Escherichia coli.
Species	Mouse
Source	E.coli
ProteinLength	120
Description	Midkine, also named MK, MK1, NEGF 2, is belonging to the neurotrophic and developmentally-regulated heparin-binding molecules family. It is encoded by the MDK gene. The Midkine protein includes five intrachain disulfide bonds which hold two domains and there are three antiparallel beta-sheets in each domain. A chondroitin sulfate proteoglycan, protein-tyrosine phosphatase zeta (PTPzeta), is a receptor for MK. MK promotes the growth, survival, and migration of various cells, and plays roles in neurogenesis and epithelial mesenchymal interactions during organogenesis. The predicted molecular weight is approximately 13.3 kDa, based on a mature peptide length of 118 amino acid residues in the mouse and 121 amino acid residues in the human. Across species, MK shows 87 % identity between the human and murine proteins.
Form	Lyophilized from a 0.2µm filtered concentrated solution in 2 × PBS, pH7.4.
Bio-activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human neutrophils is in a concentration range of 10-100 ng/ml.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 13.2 kDa, a single non-glycosylated polypeptide chain containing 120 amino acids.
AA Sequence	VAKKKEKVKKGSECSEWTWGPCTPSSKDCGMGFREGTCGAQTQRVHCKVPCNW KKEFGADCKYKFESWGACDGTGTKARQGTLLKARYNAQCQETIRVTKPCTSKTK SKTKAKKGKGKD
Endotoxin	Less than 1 EU/μg of rMuMidkine as determined by LAL method.
Purity	>96% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	Mdk
Official Symbol	Mdk
Synonyms	MDK; midkine; retinoic acid-induced differentiation factor; MK; Mek;
Gene ID	17242

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq [NM_001012335](#)

Protein Refseq [NP_001012335](#)

UniProt ID [P12025](#)

SDS-PAGE of Mdk-
567M



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA