

Recombinant Rat Il1rn protein

Cat. No. Il1rn-574R **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Rat Il1rn protein was expressed in Escherichia coli.
Species	Rat
Source	E.coli
ProteinLength	152
Description	IL-1RA was initially called the IL-1 inhibitor which is encoded by the IL1RN gene and it is a member of the interleukin 1 cytokine family. IL-1RA is secreted by various types of cells including immune cells, epithelial cells, and adipocytes. IL-RA has functions of inhibiting the activity of interleukin-1 by binding to receptor IL1R1 and preventing its association with the coreceptor IL1RAP for signaling. IL-1RA is also used in the treatment of rheumatoid arthritis, an autoimmune disease in which IL-1 plays a key role. The rat IL-1RA is a single non-glycosylated polypeptide chain containing 152 amino acids and it has been shown to block the inflammatory responses induced by IL-1 both in vitro and in vivo. Rat IL-1RA shares 89% and 73% a.a. sequence homology with murine and human IL-1RA.
Form	Lyophilized from a 0.2µm filtered concentrated solution in PBS, pH 7.4.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by inhibiting IL-1α-dependent proliferation of murine D10S cells is less than 150 ng/ml, corresponding to a specific activity of $> 6.7 \times 10^3$ IU/mg in the presence of 50 pg/ml rRtIL-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	1α.
Molecular Mass	Approximately 17.4 kDa, a single non-glycosylated polypeptide chain containing 152 amino acids.
AA Sequence	HPAGKR PCKMQAFRIWDTNQKTFYLRNNQLIAGYLQGPNTKLEEKIDMVPIDFRNVF LGIHGGKLCLSCVKSGDDTKLQLEEVNITDLNKNKEEDKRFTFIRSETGPTTSFESLA CPGWFLCTTLEADHPVSLTNTPKPCTVTKFYFQEDQ
Endotoxin	Less than 1 EU/g of rRtIL-1RA as determined by LAL method.
Purity	>96% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	Il1rn
Official Symbol	Il1rn
Synonyms	IL-1RN, IRAP, IL1 Inhibitor

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID 60582

mRNA Refseq NM_022194

Protein Refseq NP_071530

UniProt ID P25086

SDS-PAGE of Il1rn-574R

