

Recombinant Mouse Ccl5 protein

Cat. No. Ccl5-616M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse Ccl5 protein was expressed in Escherichia coli.
Species	Mouse
Source	E.coli
ProteinLength	68
Description	Chemokine (C-C motif) ligand 5 (CCL5) also known as RANTES is classified as a chemotactic cytokine or chemokine. It is encoded by the CCL5 gene in humans. CCL5 is chemotactic for T cells, eosinophils, and basophils, and plays an active role in recruiting leukocytes into inflammatory sites which has been reported to be produced by renal tubular epithelium, synovial fibroblasts and selected tumor cells. In addition, it also induces the proliferation and activation of certain natural-killer cells to form CHAK cells. Recombinant murine CCL5, a single non-glycosylated polypeptide, contains 68 amino acids and it shares 84 % and 100 % a.a. sequence identity with human and rat CCL5.
Form	Lyophilized from a 0.2µm filtered concentrated solution in 30 % Acetonitrile and 0.1 % TFA.
Bio-activity	Fully biologically active when compared to standard. The biological activity determined by a chemotaxis bioassay using human T lymphocytes is in a concentration range of 1.0-10 ng/ml.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 7.9 kDa, a single non-glycosylated polypeptide chain containing 68 amino acids.
AA Sequence	SPYGSDTTPCCFAYLSLALPRAHVKEYFYTSSKCSNLAVVFVTRRRNRQVCANPEKK WVQEYINYLEMS
Endotoxin	Less than 1 EU/g of rMuRANTES/CCL5 as determined by LAL method.
Purity	>97% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	Ccl5
Official Symbol	Ccl5
Synonyms	CCL5; chemokine (C-C motif) ligand 5; C-C motif chemokine 5; SIS-delta; small inducible cytokine A5; small-inducible cytokine A5; T-cell-specific protein RANTES; SISd; Scya5; RANTES; TCP228; MuRantes;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	20304
mRNA Refseq	NM_013653
Protein Refseq	NP_038681
UniProt ID	P30882

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA