

Active Recombinant Human TNFRSF11A protein

Cat. No. TNFRSF11A-591H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human TNFRSF11A protein was expressed in Escherichia coli.
Species	Human
Source	E.coli
ProteinLength	174
Description	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptors can interact with various TRAF family proteins, through which this receptor induces the activation of NF-kappa B and MAPK8/JNK. This receptor and its ligand are important regulators of the interaction between T cells and dendritic cells. This receptor is also an essential mediator for osteoclast and lymph node development. Mutations at this locus have been associated with familial expansile osteolysis, autosomal recessive osteopetrosis, and Paget disease of bone. Alternatively spliced transcript variants have been described for this locus.
Form	Lyophilized from a 0.2µm filtered concentrated solution in 20 mM Tris-HCl, pH 8.0, 150mM NaCl.
Bio-activity	Fully biologically active when compared to standard. The ED50 as determined by its ability to inhibit sRANK Ligand induced nuclear factor kappa B(NFkappaB) in RAW 264.7 cells is less than 50 ng/ml, corresponding to a specific activity of $> 2.0 \times 10^4$ IU/mg in the presence of 15 ng/ml of recombinant sRANK Ligand.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	Approximately 19.1 kDa, a single non-glycosylated polypeptide chain containing 174 amino acids.
AA Sequence	QIAPPCTSEKHYEHLGRCCNKCEPGKYMSSKCTTTSDSVCLPCGPDEYLDSWNEE DKCLLHKVCDTGKALVAVVAGNSTTPRRCACTAGYHWSQDCECCRRNTECAPGLG AQHPLQLNKDTVCKPCLAGYFSDAFSSTDKCRPWTNCTFLGKRVEHHGTEKSDAV CSSSLPARK
Endotoxin	Less than 0.1 EU/μg of rHusRANK Receptor as determined by LAL method.
Purity	>98% by SDS-PAGE and HPLC analysis.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. 12 months from date of receipt, -20 to -70 centigrade as supplied. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, -20 to -70 centigrade under sterile conditions after reconstitution.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤-20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	TNFRSF11A
Official Symbol	TNFRSF11A
Synonyms	TNFRSF11A; tumor necrosis factor receptor superfamily, member 11a, NFKB activator; tumor necrosis factor receptor superfamily, member 11a, activator of NFKB;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

tumor necrosis factor receptor superfamily member 11A; CD265; RANK; receptor activator of NF-KB; osteoclast differentiation factor receptor; receptor activator of nuclear factor-kappa B; loss of heterozygosity, 18, chromosomal region 1; FEO; OFE; ODFR; OST5; PDB2; OPTB7; TRANCER; LOH18CR1;

Gene ID 8792

mRNA Refseq NM_003839

Protein Refseq NP_003830

MIM 603499

UniProt ID Q9Y6Q6

**SDS-PAGE of
TNFRSF11A-591H**

