

Active Recombinant Mouse Cd86 Protein, His Tagged

Cat. No. Cd86-40M **Lot. No.** (See product label)

SPECIFICATION

Product Overview Mouse B7-2, His Tag is expressed from human 293 cells (HEK293). It contains AA Val 24-Glu 245 (Accession # NP_062261.3). This protein carries a polyhistidine tag at the C-terminus. Predicted N-terminus: Val 24

Species Mouse

Source HEK293

ProteinLength 24-245 aa

Description Cluster of Differentiation 86 (CD86) is also known as B-lymphocyte activation antigen B7-2, is a type I membrane protein that is a member of the immunoglobulin superfamily, and is constitutively expressed on interdigitating dendritic cells, Langerhans cells, peripheral blood dendritic cells, memory B cells, and germinal center B cells. Additionally, B72 is expressed at low levels on monocytes and can be upregulated through interferon γ . CD86 is the ligand for two different proteins on the T cell surface: CD28 (for autoregulation and intercellular association) and CTLA-4 (for attenuation of regulation and cellular disassociation). CD86 works in tandem with CD80 to prime T cells. Recent study has revealed that B7-2 promotes the generation of a mature APC repertoire and promotes APC function and survival. Furthermore, the B7 proteins are also involved in innate immune responses by activating NF- κ B-signaling pathway in macrophages. CD86 thus is regarded as a promising candidate for immune therapy. CD86+ macrophages in Hodgkin lymphoma patients are an independent marker for potential nonresponse to firstline-therapy.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Form	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Bio-activity	Immobilized Mouse B7-2, His Tag at 2 µg/mL (100 µL/well) can bind Rat CTLA-4, Fc Tag with a linear range of 0.2-3 ng/mL.
Molecular Mass	The protein has a calculated MW of 27.1 kDa. The protein migrates as 40-66 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 1.0 EU/µg by the LAL method/rFC method.
Purity	>95% as determined by SDS-PAGE. >95% as determined by SEC-MALS.
Storage	This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution. For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.

GENE INFORMATION

Gene Name	Cd86 CD86 antigen [<i>Mus musculus</i> (house mouse)]
Official Symbol	Cd86
Synonyms	CD86; CD86 antigen; T-lymphocyte activation antigen CD86; activation B7-2 antigen; early T cell costimulatory molecule-1; early T-cell costimulatory molecule 1; B7; B70; MB7; B7-2; B7.2; CLS1; Ly58; ETC-1; Ly-58; MB7-2; Cd28l2; TS/A-2;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA


Gene ID 12524

mRNA Refseq NM_019388

Protein Refseq NP_062261

UniProt ID P42082

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA