

Recombinant Active Mouse CXCL12 Protein, His-tagged(C-ter)

Cat. No. Cxcl12-55M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Active Mouse CXCL12 Protein with His tag (C-ter) was expressed in E. coli.
Species	Mouse
Source	E.coli
Description	This antimicrobial gene encodes a stromal cell-derived alpha chemokine member of the intercrine family. The encoded protein functions as the ligand for the G-protein coupled receptor, chemokine (C-X-C motif) receptor 4, and plays a role in many diverse cellular functions, including embryogenesis, immune surveillance, inflammation response, tissue homeostasis, and tumor growth and metastasis. Mutations in this gene are associated with resistance to human immunodeficiency virus type 1 infections. Multiple transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2014]
Form	Powder
Bio-activity	Determined by its ability to chemoattract BaF3 mouse pro-B cells transfected with human CXCR4. The ED50 for this effect is < 0.5 ng/mL.
AA Sequence	MGKPVLSYRCPCRFFESHIANVKHLKILNTPNCALQIVARLKNNNRQVCIDPKLK WIQEYLEKALNK
Endotoxin	Endotoxin level is less than 0.1 EU/μg of the protein, as determined by the LAL test.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	> 98% (by SDS-PAGE)
Applications	SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 7.4)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.

GENE INFORMATION

Gene Name	Cxcl12 chemokine (C-X-C motif) ligand 12 [<i>Mus musculus</i>]
Official Symbol	Cxcl12
Synonyms	CXCL12; chemokine (C-X-C motif) ligand 12; stromal cell-derived factor 1; C-X-C motif chemokine 12; pre-B cell growth-stimulating factor; pre-B-cell growth-stimulating factor; thymic lymphoma cell-stimulating factor; 12-O-tetradecanoylphorbol 13-acetate repressed protein 1; Pbsf; Sdf1; Tlsf; Sdf1a; Sdf1b; Tlsfa; Tlsfb; Tpar1; Scyb12; AI174028;
Gene ID	20315

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA



mRNA Refseq [NM_001012477](#)

Protein Refseq [NP_001012495](#)

SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA