

Recombinant Active Human FAS Protein (ECD), Fc-His-tagged(C-ter)

Cat. No. FAS-70H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Active Human FAS Protein (ECD) with Fc-His tag (C-ter) was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	Gln26-Asn173
Description	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor contains a death domain. It has been shown to play a central role in the physiological regulation of programmed cell death, and has been implicated in the pathogenesis of various malignancies and diseases of the immune system. The interaction of this receptor with its ligand allows the formation of a death-inducing signaling complex that includes Fas-associated death domain protein (FADD), caspase 8, and caspase 10. The autoproteolytic processing of the caspases in the complex triggers a downstream caspase cascade, and leads to apoptosis. This receptor has been also shown to activate NF-kappaB, MAPK3/ERK1, and MAPK8/JNK, and is found to be involved in transducing the proliferating signals in normal diploid fibroblast and T cells. Several alternatively spliced transcript variants have been described, some of which are candidates for nonsense-mediated mRNA decay (NMD). The isoforms lacking the transmembrane domain may negatively regulate the apoptosis mediated by the full length isoform. [provided by RefSeq, Mar

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	2011]
Form	Powder
Bio-activity	Measured by its ability to inhibit Fas Ligand-induced apoptosis of Jurkat Human acute T cell leukemia cells. The ED50 for this effect is typically 16.5-66 ng/mL in the presence of 5 ng/mL Recombinant Human Fas Ligand.
Endotoxin	Endotoxin level is less than 0.1 EU/μg of the protein, as determined by the LAL test.
Purity	> 90% (by SDS-PAGE)
Applications	Binding, FuncSt, SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C or -80°C. After reconstitution, aliquot and store at -20°C for up to one month, at 2-8°C for up to one week. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 7.4)
Reconstitution	Reconstitute to a concentration of 0.1-0.5 mg/ml in sterile distilled water.

GENE INFORMATION

Gene Name	FAS Fas (TNF receptor superfamily, member 6) [Homo sapiens]
Official Symbol	FAS

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms

FAS; Fas (TNF receptor superfamily, member 6); APT1, FAS1, TNFRSF6, tumor necrosis factor receptor superfamily, member 6; tumor necrosis factor receptor superfamily member 6; APO 1; CD95; Fas AMA; FAS 827dupA; CD95 antigen; FASLG receptor; apoptosis antigen 1; Delta Fas/APO-1/CD95; APO-1 cell surface antigen; apoptosis-mediating surface antigen FAS; tumor necrosis factor receptor superfamily, member 6; APT1; FAS1; APO-1; FASTM; ALPS1A; TNFRSF6;

Gene ID

355

mRNA Refseq

NM_000043

Protein Refseq


NP_000034

MIM

134637

UniProt ID

P25445

SDS-PAGE


Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA