

Recombinant Active Pig FGF2 Protein, His-tagged(N-ter)

Cat. No. FGF2-78P **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Active Pig FGF2 Protein with His tag (N-ter) was expressed in E. coli.
Species	Pig
Source	E.coli
Description	The protein encoded by this gene is a member of the fibroblast growth factor (FGF) family. FGF family members bind heparin and possess broad mitogenic and angiogenic activities. This protein has been implicated in diverse biological processes, such as limb and nervous system development, wound healing, and tumor growth. The mRNA for this gene contains multiple polyadenylation sites, and is alternatively translated from non-AUG (CUG) and AUG initiation codons, resulting in five different isoforms with distinct properties. The CUG-initiated isoforms are localized in the nucleus and are responsible for the intracrine effect, whereas, the AUG-initiated form is mostly cytosolic and is responsible for the paracrine and autocrine effects of this FGF. [provided by RefSeq, Jul 2008]
Form	Powder
Bio-activity	Determined by its ability to induce proliferation in 3T3 cells. The ED50 for this effect is < 2 ng/mL.
AA Sequence	AAGSITTLPALPEDGGSGAFPPGHFKDPKRLYCKNGGFFLRIHPDGRVDGVREKSD PHIKLQLQAEERGVSISIKGVCANRYLAMKEDGRLLASKCVTDECEFFERLESNNYNT YRSRKYSSWYVALKRTGQYKLGPKTGPGQKAILFLPMSAKS

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Endotoxin level is less than 0.1 EU/μg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Applications	SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 7.4) and 0.01% Sarkosyl.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.

GENE INFORMATION


Gene Name	FGF2 fibroblast growth factor 2 (basic) [<i>Sus scrofa</i>]
Official Symbol	FGF2
Synonyms	FGF2; fibroblast growth factor 2 (basic);
Gene ID	397643

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE



Pig FGF basic
recombinant protein

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA