

Recombinant Active Mouse IL12B Protein, His-tagged(C-ter)

Cat. No. IL12b-142M Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Active Mouse IL12B Protein with His tag (C-ter) was expressed in E. coli.
Species	Mouse
Source	E.coli
Description	This gene encodes a subunit of interleukin 12, a cytokine that acts on T and natural killer cells, and has a broad array of biological activities. Interleukin 12 is a disulfide-linked heterodimer composed of the 40 kD cytokine receptor like subunit encoded by this gene, and a 35 kD subunit encoded by IL12A. This cytokine is expressed by activated macrophages that serve as an essential inducer of Th1 cells development. This cytokine has been found to be important for sustaining a sufficient number of memory/effector Th1 cells to mediate long-term protection to an intracellular pathogen. Overexpression of this gene was observed in the central nervous system of patients with multiple sclerosis (MS), suggesting a role of this cytokine in the pathogenesis of the disease. The promoter polymorphism of this gene has been reported to be associated with the severity of atopic and non-atopic asthma in children. [provided by RefSeq, Jul 2008]
Form	Powder
Bio-activity	Determined by its ability to induce proliferation in T-cell enriched PBMC. The ED50 for this effect is < 0.3 ng/mL.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	MWELEKDVYVVEVDWTPDAPGETVNLTCDTPEEDDITWTSDQRHGVIGSGKTLTIT VKEFLDAGQYTCHKGGETLSHSHLLLHKKENGIWSTEILKNFKNKTFLKCEAPNYSG RFTCSWLVRNMDLKFNIKSSSSSPDSRAVTCGMASLSAEKVTLQQRDYEKYSVSC QEDVTCPTAEETLPIELALEARQQNKYENYSTSFFIRDIKPDPPKNLQMKPLKNSQV EVSWEYPDSWSTPHSYFSLKFFVRIQRKKEKMKETEEGCNQKGAFLVEKTSTEVQC KGGNVCVQAQDRYYNSSCSKWACVPCRVR
Endotoxin	Endotoxin level is less than 0.1 EU/μg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Applications	SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 7.4)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.

GENE INFORMATION

Gene Name	Il12b interleukin 12b [<i>Mus musculus</i>]
Official Symbol	Il12b

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms IL12B; interleukin 12b; interleukin-12 subunit beta; CLMF p40; IL-12 p40; IL-12 subunit p40; IL-23 subunit p40; cytotoxic lymphocyte maturation factor 40 kDa subunit; p40; Il-12b; Il12p40; Il-12p40;

Gene ID [16160](#)

mRNA Refseq [NM_008352](#)

Protein Refseq [NP_032378](#)

SDS-PAGE

