

Recombinant Active Mouse IL18 Protein, His-tagged(C-ter)

Cat. No. IL18-155M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Active Mouse IL18 Protein with His tag (C-ter) was expressed in E. coli.
Species	Mouse
Source	E.coli
Description	The protein encoded by this gene is a proinflammatory cytokine that augments natural killer cell activity in spleen cells, and stimulates interferon gamma production in T-helper type I cells. Alternatively spliced transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Aug 2011]
Form	Powder
Bio-activity	Determined by its ability to induce IFN gamma secretion in KG-1 cells. The ED50 for this effect is < 0.5 µg/mL.
AA Sequence	MNFGRLHCTTAVIRNINDQVLFVDKRQPVFEDMTDIDQSASEPQTRLIIYMYKDSEVR GLAVTL SVKDSKMSTLSCKNKIISFEEMDPPENIDDIQSDLIFFQKRVP GHNKMEFES SLYEGHFLACQKEDDAFKLILKKKDENGDKSVMFTLTNLHQS
Endotoxin	Endotoxin level is less than 0.1 EU/µg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Applications	SDS-PAGE

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 8.0)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 µg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.
GENE INFORMATION	
Gene Name	IL18 interleukin 18 [<i>Mus musculus</i>]
Official Symbol	IL18
Synonyms	IL18; interleukin 18; interleukin-18; IL-1 gamma; interleukin-1 gamma; IFN-gamma-inducing factor; interferon gamma-inducing factor; interferon-gamma-inducing factor; Igif; IL-18;
Gene ID	16173
mRNA Refseq	NM_008360
Protein Refseq	NP_032386