

Recombinant Active Pig IL4 Protein, His-tagged(C-ter)

Cat. No. IL4-205P **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Active Pig IL4 Protein with His tag (C-ter) was expressed in E. coli.
Species	Pig
Source	E.coli
Description	The protein encoded by this gene is a pleiotropic cytokine produced by activated T cells. This cytokine is a ligand for interleukin 4 receptor. The interleukin 4 receptor also binds to IL13, which may contribute to many overlapping functions of this cytokine and IL13. STAT6, a signal transducer and activator of transcription, has been shown to play a central role in mediating the immune regulatory signal of this cytokine. This gene, IL3, IL5, IL13, and CSF2 form a cytokine gene cluster on chromosome 5q, with this gene particularly close to IL13. This gene, IL13 and IL5 are found to be regulated coordinately by several long-range regulatory elements in an over 120 kilobase range on the chromosome. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported. [provided by RefSeq, Jul 2008]
Form	Powder
Bio-activity	Determined by its ability to induce proliferation in TF-1 cells. The ED50 for this effect is < 0.6 ng/mL.
AA Sequence	MHKCDITLQEIIKTLNILTARKNSCMELPVTDVFAAPENTTEKETFCRASTVLRHIYRH HTCMKSLLSGLDRNLSSMANMTCSVHEAKKSTLKDFLERLKTIMKEKYSKC

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Endotoxin	Endotoxin level is less than 0.1 EU/μg of the protein, as determined by the LAL test.
Purity	> 98% (by SDS-PAGE)
Applications	SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 7.4)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.

GENE INFORMATION

Gene Name	IL4 interleukin 4 [Sus scrofa]
Official Symbol	IL4
Synonyms	IL4; interleukin 4; interleukin-4; IL-4; BSF-1; B-cell stimulatory factor 1; lymphocyte stimulatory factor 1;
Gene ID	397225
mRNA Refseq	NM_214123


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq

NP_999288

SDS-PAGE Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA