

Recombinant Active Human INHBA Protein, His-tagged(C-ter)

Cat. No. INHBA-220H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Active Human INHBA Protein with His tag (C-ter) was expressed in E. coli.
Species	Human
Source	E.coli
Description	The inhibin beta A subunit joins the alpha subunit to form a pituitary FSH secretion inhibitor. Inhibin has been shown to regulate gonadal stromal cell proliferation negatively and to have tumor-suppressor activity. In addition, serum levels of inhibin have been shown to reflect the size of granulosa-cell tumors and can therefore be used as a marker for primary as well as recurrent disease. Because expression in gonadal and various extragonadal tissues may vary severalfold in a tissue-specific fashion, it is proposed that inhibin may be both a growth/differentiation factor and a hormone. Furthermore, the beta A subunit forms a homodimer, activin A, and also joins with a beta B subunit to form a heterodimer, activin AB, both of which stimulate FSH secretion. Finally, it has been shown that the beta A subunit mRNA is identical to the erythroid differentiation factor subunit mRNA and that only one gene for this mRNA exists in the human genome. [provided by RefSeq, Jul 2008]
Form	Powder
Bio-activity	Determined by its ability to induce hemoglobin expression in K562 cells. The ED50 for this effect is < 0.85 ng/mL. The specific activity of recombinant human IGF-I is

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	approximately $> 1.4 \times 10^3$ IU/mg.
AA Sequence	MGLECDGKVNICKKKQFFVSFKDIGWNDWIIAPSGYHANYCEGECPSHIAGTSGSSL SFHSTVINHYRMRGHSPFANLKSCCVPTKLRPMSMLYYDDGQNIKKDIQNMIVEEC GCS
Endotoxin	Endotoxin level is less than 0.1 EU/ μ g of the protein, as determined by the LAL test.
Purity	> 95% (by SDS-PAGE)
Applications	SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 8.0) and 0.1% Sarkosyl.
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μ g/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.


GENE INFORMATION

Gene Name	INHBA inhibin, beta A [Homo sapiens]
Official Symbol	INHBA

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Synonyms	INHBA; inhibin, beta A; inhibin, beta A (activin A, activin AB alpha polypeptide); inhibin beta A chain; Inhibin, beta-1; activin beta-A chain; FSH-releasing protein; erythroid differentiation factor; erythroid differentiation protein; follicle-stimulating hormone-releasing protein; EDF; FRP;
Gene ID	3624
mRNA Refseq	NM_002192
Protein Refseq	NP_002183
MIM	147290
UniProt ID	P08476
SDS-PAGE	 Human Activin A recombinant protein