

Recombinant Active Mouse INHBB Protein, His-tagged(C-ter)

Cat. No. Inhbb-221M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Active Mouse INHBB Protein with His tag (C-ter) was expressed in E. coli.
Species	Mouse
Source	E.coli
Description	The inhibin beta B subunit joins the alpha subunit to form a pituitary FSH secretion inhibitor. Inhibin has been shown to regulate gonadal stromal cell proliferation negatively and to have tumour-suppressor activity. In addition, serum levels of inhibin have been shown to reflect the size of granulosa-cell tumors and can therefore be used as a marker for primary as well as recurrent disease. Because expression in gonadal and various extragonadal tissues may vary severalfold in a tissue-specific fashion, it is proposed that inhibin may be both a growth/differentiation factor and a hormone. Furthermore, the beta B subunit forms a homodimer, activin B, and also joins with the beta A subunit to form a heterodimer, activin AB, both of which stimulate FSH secretion. [provided by RefSeq, Jul 2008]
Form	Powder
Bio-activity	Determined by its ability to induce hemoglobin expression in K562 cells. The ED50 for this effect is < 1 ng/mL.
AA Sequence	MGLECDGRTSLCCRQQFFIDFRLIGWNDWIIAPTGYGNYCEGSCPAYLAGVPGSA SSFHTAVVNQYRMRGLNPGPVNSCCIPTKLSSMSMLYFDDEYNIVKRDVPNMIVVEE

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

CGCA	
Endotoxin	Endotoxin level is less than 0.1 EU/μg of the protein, as determined by the LAL test.
Purity	> 95% (by SDS-PAGE)
Applications	SDS-PAGE
Notes	For laboratory research only, not for drug, diagnostic or other use.
Storage	Lyophilized protein should be stored at -20°C. After reconstitution, aliquot and store at -20°C or -80°C for up to one month. Storage in frost free freezers is not recommended. Avoid repeated freeze/thaw cycles. Suggest spin the vial prior to opening.
Storage Buffer	PBS (pH 8.0)
Reconstitution	It is recommended to reconstitute the lyophilized protein in sterile water to a concentration not less than 100 μg/mL and incubate the stock solution for at least 20 min at room temperature to make sure the protein is dissolved completely.
GENE INFORMATION	
Gene Name	Inhbb inhibin beta-B [Mus musculus]
Official Symbol	Inhbb
Synonyms	INHBB; inhibin beta-B; inhibin beta B chain; activin beta-B chain;
Gene ID	16324

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq [NM_008381](#)

Protein Refseq [NP_032407](#)

SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA