

Recombinant Human/Bovine/Porcine TGFB1 Protein

Cat. No. TGFB1-37H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Recombinant human/bovine/porcine TGF- β 1 PLUS protein without tag was expressed in E. coli.

Animal-free (AOF) and carrier protein-free.

Mass spectrometry: single species with expected mass.

Analytical reversed-phase: single sharp peak.

Recovery from stock vial: >95%

Species

Human

Source

E.coli

Description

This gene encodes a secreted ligand of the TGF-beta (transforming growth factor-beta) superfamily of proteins. Ligands of this family bind various TGF-beta receptors leading to recruitment and activation of SMAD family transcription factors that regulate gene expression. The encoded preproprotein is proteolytically processed to generate a latency-associated peptide (LAP) and a mature peptide, and is found in either a latent form composed of a mature peptide homodimer, a LAP homodimer, and a latent TGF-beta binding protein, or in an active form consisting solely of the mature peptide homodimer. The mature peptide may also form heterodimers with other TGFB family members. This encoded protein regulates cell proliferation, differentiation and growth, and can modulate expression and activation of other growth factors including interferon gamma and tumor necrosis factor alpha. This gene is frequently upregulated in tumor cells, and mutations in this gene result in Camurati-Engelmann disease.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	24 kDa (dimer)
Endotoxin	<0.005 EU/μg protein (below level of detection)
Applications	Maintenance of pluripotency in human iPSC; cell culture media optimisation and replacement of insulin in defined media
Storage	Resuspend in 10mM HCl at >100 μg/mL, prepare single use aliquots, add carrier protein if desired and store frozen at -20 or -80 centigrade
Storage Buffer	Lyophilized from acetonitrile, TFA
GENE INFORMATION	
Gene Name	TGFB1 transforming growth factor beta 1 [Homo sapiens (human)]
Official Symbol	TGFB1
Synonyms	TGFB1; transforming growth factor beta 1; CED; LAP; DPD1; TGFB; IBDIMDE; TGFbeta; TGF-beta1; transforming growth factor beta-1 proprotein; TGF-beta-1; latency-associated peptide; prepro-transforming growth factor beta-1; transforming growth factor beta1
Gene ID	7040
mRNA Refseq	NM_000660
Protein Refseq	NP_000651
MIM	190180


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID
P01137
Bioactivity

TGF- β 1 PLUS is highly bioactive compared to mammalian-expressed TGF- β 1 preparations from other suppliers. Comparative activity determined using a quantitative luciferase reporter assay in transiently transfected HEK293T cells. Cells were treated with a serial dilution of TGF- β 1. Firefly luciferase activity is measured and normalized to the control Renilla luciferase activity. TGF- β 1 PLUS EC₅₀ 1.5 pM (0.037 ng/mL). Suppliers 1 and 2 have EC₅₀ ~3.5 pM and 38 pM respectively.

Purity


TGF- β 1 PLUS dimer migrates as a single band at 24 kDa in non-reducing and 13 kDa as a single monomeric species upon reduction (R). High purity yield of dimeric protein (bioactive form). Purified recombinant protein was resolved using 15% w/v SDS-PAGE in reduced and non-reduced conditions and stained with Coomassie Brilliant Blue R250.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA