

Recombinant Human CCL2, Fc-tagged

Cat. No. CCL2-130H Lot. No. (See product label)

SPECIFICATION

Product Overview	The extracellular domain of human CCL2 (aa 24-99) is fused to the N-terminus of the Fc region of human IgG1.
Species	Human
Source	CHO
ProteinLength	24-99 a.a.
Description	CCL2, also called monocyte chemotactic protein-1 (MCP-1), is a member of the C-C or β chemokine family that is best known as a chemotactic agent for mononuclear cells. Fibroblasts, glioma cells, smooth muscle cells, endothelial cells, lymphocytes and mononuclear phagocytes can produce CCL2 either constitutively or upon mitogenic stimulation, but monocytes and macrophages appear to be the major source. In addition to its chemotactic activity, CCL2 induces enzyme and cytokine release by monocytes, NK cells and lymphocytes, and histamine release by basophils that express its receptor CCR2. Additionally, it promotes Th2 polarization in CD4⁺ T cells. CCL2-mediated recruitment of monocytes to sites of inflammation is proposed to play a role in the pathology of atherosclerosis, multiple sclerosis and allergic asthma.
Form	Lyophilized from 0.2 μ m-filtered solution in PBS.
Purity	$\geq$ 98% (SDS-PAGE)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Stability	Stable for at least 1 year after receipt when stored at -20°C. Working aliquots are stable for up to 3 months when stored at -20°C.
Storage	Short Term Storage: +4°C; Long Term Storage: -20°C. Avoid freeze/thaw cycles.
GENE INFORMATION	
Gene Name	CCL2 chemokine (C-C motif) ligand 2 [Homo sapiens]
Official Symbol	CCL2
Synonyms	CCL2; chemokine (C-C motif) ligand 2; SCYA2, small inducible cytokine A2 (monocyte chemotactic protein 1, homologous to mouse Sig je); C-C motif chemokine 2; GDCF 2; HC11; MCAF; MCP 1; MCP1; MGC9434; monocyte chemoattractant protein 1; monocyte chemotactic and activating factor; monocyte chemotactic protein 1; homologous to mouse Sig je; monocyte secretory protein JE; small inducible cytokine subfamily A (Cys Cys); member 2; SMC CF; small-inducible cytokine A2; monocyte chemoattractant protein-1; small inducible cytokine subfamily A (Cys-Cys), member 2; small inducible cytokine A2 (monocyte chemotactic protein 1, homologous to mouse Sig-je); MCP-1; SCYA2; GDCF-2; SMC-CF; HSMCR30;
Gene ID	6347
mRNA Refseq	NM_002982
Protein Refseq	NP_002973
UniProt ID	P13500
Chromosome Location	17q11.2-q21.1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway

Activation of Genes by ATF4, organism-specific biosystem; Chagas disease (American trypanosomiasis), organism-specific biosystem; Chagas disease (American trypanosomiasis), conserved biosystem; Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem;

Function

CCR2 chemokine receptor binding; CCR2 chemokine receptor binding; G-protein coupled receptor binding; chemokine activity; heparin binding; protein kinase activity; receptor binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA