

Recombinant Human CD86 Protein, C-His-tagged

Cat. No. CD86-186H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human CD86 Protein with C-His tag was expressed in E. coli.
Species	Human
Source	E.coli
Description	CD80 (B7-1, BB1) and CD86 (B7-2, B70) are members of the B7 family of cell surface ligands that regulate T cell activation and immune responses. CD80 is expressed on activated antigen presenting cells, including dendritic cells, B cells, monocytes, and macrophages. CD86 is expressed on resting monocytes, dendritic cells, activated B lymphocytes, and can be further upregulated in the presence of inflammation. CD80 and CD86 are ligands for CD28, which functions as a T cell costimulatory receptor. Interaction of CD28 with CD80 or CD86 provides the second signal required for naïve T cell activation, T cell proliferation, and acquisition of effector functions. Alternatively, CD80 and CD86 also act as ligands to CTLA-4, which results in the downregulation of T cell activity.
Molecular Mass	~25 kDa
AA Sequence	APLKIQAYFNETADLPCQFANSQNQSLSELVFWQDQENLVLNEVYLGKEKFDSVH SKYMGRTSFDSDSWTLRLHNLQIKDKGLYQCIIHHKKPTGMIRIHQMNSELSVLNFS QPEIVPISNITENVYINLTCSSIHGYPEPKKMSVLLRTKNSTIEYDGMQKSQDNVTEL YDVSISLSVSFPDVTSNMTIFCILETDKTRLLSSPFSIELEDPQPPPDHIP
Purity	Transferred into competent cells and the supernatant was purified by NI column

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

affinity chromatography and the purity is > 85% (by SDS-PAGE).

Notes

For research use only, not for use in diagnostic procedure.

Storage

Store at 4 centigrade short term. Aliquot and store at -20 centigrade long term. Avoid freeze-thaw cycles.

Concentration

≥0.5 mg/mL

Storage Buffer

PBS, 4M Urea, pH7.4

GENE INFORMATION

Gene Name

CD86 CD86 molecule [Homo sapiens (human)]

Official Symbol

CD86

Synonyms

CD86; CD86 molecule; CD28LG2, CD86 antigen (CD28 antigen ligand 2, B7 2 antigen); T-lymphocyte activation antigen CD86; B lymphocyte antigen B7 2; B7 2; B7.2; BU63; FUN-1; CTLA-4 counter-receptor B7.2; B-lymphocyte activation antigen B7-2; CD86 antigen (CD28 antigen ligand 2, B7-2 antigen); B70; B7-2; LAB72; CD28LG2; MGC34413;

Gene ID

942

mRNA Refseq

NM_001206924

Protein Refseq

NP_001193853

MIM

601020


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID P42081

SDS-PAGE



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA