

Recombinant Human CD72 Protein, C-His-tagged

Cat. No. CD72-207H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human CD72 Protein with C-His tag was expressed in E. coli.
Species	Human
Source	E.coli
Description	CD5 has been identified as a transmembrane glycoprotein that is expressed on 70% of normal peripheral blood lymphocytes and on virtually all T lymphocytes in thymus and peripheral blood. Activation of T cells through the T cell receptor (TCR) results in tyrosine phosphorylation of CD5, and the absence of CD5 renders T cells hyper-responsive to TCR-mediated activation. CD5 associates with the TCR/CD3 ζ chain, and with the Src family kinase, Lck p56. The C-type lectin superfamily member CD72 is a cell surface negative regulator of B cell activation from the pro-B through the mature B cell stage. CD72 serves as a receptor for CD5. The ability of lymphocytes to respond to antigenic or mitogenic stimulation utilizes both positive and negative regulatory proteins that influence the threshold for responsiveness. The human CD72 gene maps to chromosome 9p13.3 and encodes a transmembrane glycoprotein that contains an immunoreceptor tyrosine-based inhibition motif (ITIM). Upon tyrosine phosphorylation, the CD72 ITIM recruits SH2-containing phosphatases such as SHP-1, resulting in downregulation of cell activation. CD72^{-/-} mice contain hyperproliferative B cells.
Molecular Mass	~27 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	RYLQVSQQLQQTNRVLEVTNSSLRQQRLKITQLGQSAEDLQGSRRRLAQSQEALQ VEQRAHQAAEGQLQACQADRQKTKETLQSEEQRRRALEQKLSNMENRLKPFFTCG SADTCCPSGWIMHQKSCFYISLTSKNWQESQKQCETLSSKLATFSEIYPQSHSYFL NSLLPNGGSGNSYWTGLSSNKDWKLTDDTQRTRTYAQSSKCNKVHKTWSWWTLE SESCRSSLPYICEMTAFRFPD
Purity	Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).
Notes	For research use only, not for use in diagnostic procedure.
Storage	Store at 4 centigrade short term. Aliquot and store at -20 centigrade long term. Avoid freeze-thaw cycles.
Concentration	≥0.5 mg/mL
Storage Buffer	PBS, 4M Urea, pH7.4

GENE INFORMATION

Gene Name	CD72 CD72 molecule [Homo sapiens (human)]
Official Symbol	CD72
Synonyms	CD72; CD72 molecule; CD72 antigen; B-cell differentiation antigen CD72; CD72b; LYB2; lyb-2;
Gene ID	971
mRNA Refseq	NM_001782

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq NP_001773

MIM 107272

UniProt ID P21854

SDS-PAGE



 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA