

Active Recombinant Human TNFRSF4, Fc-tagged

Cat. No. TNFRSF4-441H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human OX40 (aa 29-214) is fused at the C-terminus to the Fc portion of human IgG1.
Species	Human
Source	HEK293
ProteinLength	29-214 a.a.
Description	OX40 binds to receptors on T-cells, preventing them from dying and subsequently increasing cytokine production. OX40 has a critical role in the maintenance of an immune response. OX40:Fc was shown to prevent OX40 from reaching the T-cell receptors, thus reducing the T-cell response. Experiments in mice have demonstrated that OX40:Fc can reduce the symptoms associated with the cytokine storm (an immune overreaction) while allowing the immune system to fight off the virus successfully.
Form	Lyophilized. Contains PBS.
Bio-activity	Blocks T cell proliferation induced by CD134L.
Molecular Mass	~70kDa (SDS-PAGE)
Purity	≥95% (SDS-PAGE)
Stability	Stable for at least 6 months after receipt when stored at -20°C.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	Short Term Storage: +4°C; Long Term Storage: -20°C. After reconstitution, prepare aliquots and store at -20°C. Avoid freeze/thaw cycles. PBS containing at least 0.1% BSA should be used for further dilutions.
Concentration	1mg/ml after reconstitution.
GENE INFORMATION	
Gene Name	TNFRSF4 tumor necrosis factor receptor superfamily, member 4 [Homo sapiens]
Official Symbol	TNFRSF4
Synonyms	TNFRSF4; tumor necrosis factor receptor superfamily, member 4; TXGP1L; tumor necrosis factor receptor superfamily member 4; ACT35; CD134; OX40; OX40 antigen; ACT35 antigen; ATC35 antigen; CD134 antigen; OX40 homologue; OX40L receptor; OX40 cell surface antigen; lymphoid activation antigen ACT35; TAX transcriptionally-activated glycoprotein 1 receptor; tax-transcriptionally activated glycoprotein 1 receptor;
Gene ID	7293
mRNA Refseq	NM_003327
Protein Refseq	NP_003318
MIM	600315
UniProt ID	P43489
Chromosome Location	1p36

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Downstream signaling in naive CD8+ T cells, organism-specific biosystem;
Function	binding; receptor activity; tumor necrosis factor-activated receptor activity;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA