

Recombinant Human Fms-related Tyrosine Kinase 4, His-tagged

Cat. No. FLT4-161H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human FLT4 produced in <i>Insect cells</i> fused with a carboxy-terminal 6X his-tag. The recombinant mature sVEGFR-3/FLT-4 is a glycosylated monomeric protein. The sVEGFR-3/FLT-4 monomers have a mass of approximately 120kDa. The soluble receptor protein consists of all 7 extracellular domains (Met1-Glu774).
Species	Human
Source	Insect Cells
Protein Length	1-774 a.a.
Description	The FLT-4 cDNA encodes a 1298 amino acid (aa) residue precursor protein with a 23 aa residue signal peptide. Mature VEGFR-3/FLT-4 is composed of a 751 aa residue extracellular domain, a 22 aa transmembrane domain and a 482 aa residue cytoplasmic domain. Both VEGF family members VEGF-C and VEGF-D have been shown to bind and activate VEGFR-3/FLT-4. The Flt-4 gene is widely expressed in the early embryo but becomes restricted to the lymphatic endothelial a latter stages of development. It is important for lymphangiogenesis. All three VEGF receptors belong to the class III subfamily of receptor tyrosine kinases (RTKs) characterised by the seven immunoglobulin like loops in the extracellular domain. The expression of VEGFR-1 to -3 is almost exclusively restricted to hematopoietic precursor cells, vascular and lymphatic endothelial cells and to the monocyte/macrophage lineage. They play key roles in vasculogenesis, hematopoiesis, angiogenesis and

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	lymphangiogenesis.
Subunit	Monomeric glycoprotein.
Purity	>90%, by SDS-PAGE and visualised by silver stain.
Endotoxin Level	<0.1ng per µg of sVEGFR-3.
Buffer	PBS.
Formulation	Lyophilized.
Biologically Activity	Measured by its ability to bind recombinant rat VEGF-C in a functional solid phase binding assay. Immobilised recombinant human sVEGFR-3/FLT-4 at 5µg/ml can bind recombinant rat VEGF-C in a linear range of 8-500ng/ml.
Reconstitution	The lyophilised sVEGFR-3/FLT-4 is soluble in water and most aqueous buffers. The lyophilised sVEGFR-3/FLT-4 should be reconstituted in PBS or medium to a concentration not lower than 100µg/ml.
Stability	Lyophilised samples are stable for greater than six months at -20°C to -70°C. Reconstituted sVEGFR-3/FLT-4 should be stored in working aliquots at -20°C. Avoid repeated freeze-thaw cycles!

GENE INFORMATION

Gene Name	FLT4 fms-related tyrosine kinase 4 [Homo sapiens]
Synonyms	FLT4; fms-related tyrosine kinase 4; PCL; FLT41; LMPH1A; VEGFR3; soluble VEGFR3 variant 1; soluble VEGFR3 variant 2; soluble VEGFR3 variant 3; vascular endothelial growth factor receptor 3; EC 2.7.10.1; 3` partial; Tyrosine-protein kinase

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

receptor FLT4; VEGFR-3	
Gene ID	2324
mRNA Refseq	NM_002020
Protein Refseq	NP_002011
MIM	136352
UniProt ID	P35916
Chromosome Location	5q34-q35
Pathway	Cytokine-cytokine receptor interaction; Focal adhesion; Signaling by VEGF
Function	ATP binding; growth factor binding; nucleotide binding; receptor activity; transferase activity; vascular endothelial growth factor receptor activity

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA