

Active Recombinant Human GTF2A1, P55, GST-tagged

Cat. No. GTF2A1-1173H **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Species Human

Source E.coli

Description

The Transcription Factor IIA (TFIIA) has been shown to bind to the TBP-DNA complex and to increase the affinity of TBP for the TATA element. Human TFIIA consists of three subunits of 35 kDa (α -subunit), 19 kDa (β -subunit) and 12 kDa (γ -subunit). The α - and β -subunits are derived from the product, p55, of a single gene by an unknown mechanism. However, recombinant p55, in combination with the 12 kDa subunit, retains native TFIIA activity. GST-p55 subunit of TFIIA is isolated from a strain of *E.coli* that contains the coding sequence of human p55 of TFIIA under the control of T7 promoter. GST-TFIIA can be used to study protein-protein interactions. Protein is greater than 95% homogeneous and contains no detectable proteases, DNase and RNase activity.

Usage For in vitro use only.

Form Liquid. Supplied in 20 mM Tris-HCl pH 8.0, 100 mM KCl, 0.2 mM EDTA, 1 mM DTT, 20% glycerol.

Activity 100 ng are sufficient for a protein-protein interaction assay.

Purity > 95% by SDS-PAGE.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage Quality guaranteed for 12 months, store at -80°C. Avoid freeze / thaw cycles.

GENE INFORMATION

Gene Name [GTF2A1 general transcription factor IIA, 1, 19/37kDa \[Homo sapiens \]](#)

Synonyms GTF2A1; general transcription factor IIA, 1, 19/37kDa; TF2A1; TFIIA; MGC129969; MGC129970; TFIIA alpha, p55, isoform 1; glucose regulated protein, 58kD pseudogene; TFIIA-42; Transcription initiation factor IIA subunit 1; Transcription initiation factor IIA alpha chain; TFIIA p35 subunit; TFIIA p19 subunit

Gene ID [2957](#)

mRNA Refseq [NM_015859](#)

Protein Refseq [NP_056943](#)

MIM [600520](#)

UniProt ID [P52655](#)

Chromosome Location 14q31

Pathway Basal transcription factors; Gene Expression; HIV Infection; Transcription


Function DNA binding; TATA-binding protein binding; general RNA polymerase II transcription factor activity; protein heterodimerization activity; transcription coactivator activity; transcription factor binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

PDB rendering based
on 1nvp.



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA