

Recombinant Full Length Human UCHL5 Protein, C-Flag-tagged

Cat. No. UCHL5-2070HFL **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Full Length Human UCHL5 Protein, fused to Flag-tag at C-terminus, was expressed in Mammalian cells.
Species	Human
Source	Mammalian Cells
Description	Enables endopeptidase inhibitor activity; proteasome binding activity; and thiol-dependent deubiquitinase. Involved in negative regulation of proteasomal ubiquitin-dependent protein catabolic process; positive regulation of smoothened signaling pathway; and protein deubiquitination. Located in cytosol; nucleolus; and nucleoplasm. Colocalizes with Ino80 complex.
Form	25 mM Tris HCl, pH 7.3, 100 mM glycine, 10% glycerol.
Molecular Mass	37.4 kDa
AA Sequence	MTGNAGEWCLMESDPGVFTELIKGFGRGAQVEEIWSLEPENFEKLKPVHGLIFLFLK WQPGEEPAGSVVQ DSRLDTIFFAKQVINNACATQAIVSVLLNCTHQDVHLGETLSEF KEFSQSFDAAMKGLALSNSDVIRQVH NSFARQQMFEFDTKTSAKEEDAFHFVSYVP VNGRLYELDGLREGPIDLGACNQDDWISAVRPVIEKRIQK YSEGEIRFNLMAIVSDR KMIYEQKIAELQRQLAEEEPMDTDQGNSMLSAIQSEVAKNQMLIEEEVQKLR YKIE NIRRKHNYLPFIMELLKTLAEHQQLIPLVEKAKEKQNAKKAQETK TRTRPLEQKLISE EDLAANDILDYKDDDDKV

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	> 80% as determined by SDS-PAGE and Coomassie blue staining.
Stability	Stable for 12 months from the date of receipt of the product under proper storage and handling conditions. Avoid repeated freeze-thaw cycles.
Storage	Store at -80 centigrade.
Concentration	>50 ug/mL as determined by microplate BCA method.
Preparation	Recombinant protein was captured through anti-DDK affinity column followed by conventional chromatography steps.
Protein Families	Druggable Genome, Protease
Full Length	Full L.

GENE INFORMATION

Gene Name	UCHL5 ubiquitin C-terminal hydrolase L5 [Homo sapiens (human)]
Official Symbol	UCHL5
Synonyms	Enables endopeptidase inhibitor activity; proteasome binding activity; and thiol-dependent deubiquitinase. Involved in negative regulation of proteasomal ubiquitin-dependent protein catabolic process; positive regulation of smoothened signaling pathway; and protein deubiquitination. Located in cytosol; nucleolus; and nucleoplasm. Colocalizes with Ino80 complex.
Gene ID	51377
mRNA Refseq	NM_015984.5

 Tel: 1-631-559-9269 1-516-512-3133


 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Protein Refseq NP_057068.1

MIM 610667

UniProt ID Q9Y5K5



Coomassie blue staining of purified UCHL5 protein.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA