

Recombinant Full Length Rat Voltage-Dependent Calcium Channel Gamma-8 Subunit(Cacng8) Protein, His-Tagged

Cat. No. RFL10314RF **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Full Length Rat Voltage-dependent calcium channel gamma-8 subunit(Cacng8) Protein (Q8VHW5) (1-421aa), fused to N-terminal His tag, was expressed in E. coli.
Species	Rat
Source	E.coli
ProteinLength	Full Length (1-421)
Form	Lyophilized powder
AA Sequence	MESLKRWNEERGLWCEKGVQVLLTTIGAFAAFGMLMTIAISTDYWLYTRALICNTTNLT AG DDGPPHRGGSGSSEKKDPGGLTHSGLWRICCLEGLKRGVCVKINHFPEDTDYD HDSAEYL LRVVRASSIFPILSAILLLLGGVCVAASRVYKSKRNIILGAGILFVAAGLSNII GVIVYI SANAGEPGPKRDEEKKNHYSYGWSFYFGGLSFILAEVIGVLAVNIYIERSRE AHCQSRSD LLKAGGGAGGSGGSGPSAILRLPSYRFRYRRRSRSSSRGSSEASPSR DASPGGPGGPGFA STDISMYTLSRDPSKGSVAAGLASAGGGGGGAGVGAYGGAA GAAGGGGTGSEDRGSSAG FLTLHNAFPKEAASGVTVTVTGPPAAPAPAPPAPAA PAPGTLSKEAAASNTNTLNKTKTP V
Purity	Greater than 90% as determined by SDS-PAGE.
Notes	Repeated freezing and thawing is not recommended. Store working aliquots at 4°C

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	for up to one week.
Storage	Store at -20°C/-80°C upon receipt, aliquoting is necessary for mutiple use. Avoid repeated freeze-thaw cycles.
Storage Buffer	Tris/PBS-based buffer, 6% Trehalose, pH 8.0
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Please reconstitute protein in deionized sterile water to a concentration of 0.1-1.0 mg/mL. We recommend to add 5-50% of glycerol (final concentration) and aliquot for long-term storage at -20°C/-80°C. Our default final concentration of glycerol is 50%. Customers could use it as reference.
GENE INFORMATION	
Gene Name	Cacng8
Synonyms	Cacng8; Voltage-dependent calcium channel gamma-8 subunit; Neuronal voltage-gated calcium channel gamma-8 subunit; Transmembrane AMPAR regulatory protein gamma-8; TARP gamma-8
UniProt ID	Q8VHW5