

Active Recombinant Human CD19 Protein, His Tagged

Cat. No. CD19-26H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Human CD19 (20-291) Protein, His Tag is expressed from CHO cells. It contains AA Pro 20-Lys 291 (Accession # P15391-1). This protein carries a polyhistidine tag at the C-terminus. Predicted N-terminus: Pro 20
Species	Human
Source	CHO
ProteinLength	20-291 aa
Description	B-lymphocyte antigen CD19 is also known as CD19 (Cluster of Differentiation 19), is a single-pass type I membrane protein which contains two Ig-like C2-type (immunoglobulin-like) domains. CD19 is expressed on follicular dendritic cells and B cells. In fact, it is present on B cells from earliest recognizable B-lineage cells during development to B-cell blasts but is lost on maturation to plasma cells. It primarily acts as a B cell co-receptor in conjunction with CD21 and CD81. Upon activation, the cytoplasmic tail of CD19 becomes phosphorylated, which leads to binding by Src-family kinases and recruitment of PI-3 kinase. As on T cells, several surface molecules form the antigen receptor and form a complex on B lymphocytes. The (almost) B cell-specific CD19 phosphoglycoprotein is one of these molecules. The others are CD21 and CD81. These surface immunoglobulin (sIg)-associated molecules facilitate signal transduction. On living B cells, anti-immunoglobulin antibody mimicking exogenous antigen causes CD19 to bind to sIg and internalize with it. The reverse process has not been demonstrated, suggesting that formation of

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

this receptor complex is antigen-induced. This molecular association has been confirmed by chemical studies. Mutations in CD19 are associated with severe immunodeficiency syndromes characterized by diminished antibody production. CD19 has been shown to interact with: CD81, CD82, Complement receptor 2, and VAV2.

Form	Lyophilized from 0.22 µm filtered solution in PBS, pH7.4 with trehalose as protectant.
Bio-activity	Immobilized Human CD19 (20-291) Protein, His Tag at 2 µg/mL (100 µL/well) can bind Monoclonal Anti-Human CD19 Antibody, Mouse IgG2a with a linear range of 0.1-3 ng/mL.
Molecular Mass	The protein has a calculated MW of 32.0 kDa. The protein migrates as 45-60 kDa when calibrated against Star Ribbon Pre-stained Protein Marker under reducing (R) condition (SDS-PAGE) due to glycosylation.
Endotoxin	Less than 1.0 EU/µg by the LAL method/rFC method.
Purity	>90% as determined by SDS-PAGE.
Storage	This product is stable after storage at: -20 to -70 centigrade for 12 months in lyophilized state; -70 centigrade for 3 months under sterile conditions after reconstitution. For long term storage, the product should be stored at lyophilized state at -20 centigrade or lower. Please avoid repeated freeze-thaw cycles.
Reconstitution	It is recommended that sterile water be added to the vial to prepare a stock solution of 0.2 µg/µL. Centrifuge the vial at 4 centigrade before opening to recover the entire contents.

GENE INFORMATION

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	CD19 CD19 molecule [Homo sapiens (human)]
Official Symbol	CD19
Synonyms	CD19; CD19 molecule; CD19 antigen; B-lymphocyte antigen CD19; differentiation antigen CD19; T-cell surface antigen Leu-12; B-lymphocyte surface antigen B4; B4; CVID3; MGC12802;
Gene ID	930
mRNA Refseq	NM_001178098
Protein Refseq	NP_001171569
MIM	107265
UniProt ID	P15391

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA