

Recombinant Human VEGFA Protein

Cat. No. VEGFA-0062H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human VEGFA Protein without tag was expressed in HEK293.
Species	Human
Source	HEK293
Description	This gene is a member of the PDGF/VEGF growth factor family. It encodes a heparin-binding protein, which exists as a disulfide-linked homodimer. This growth factor induces proliferation and migration of vascular endothelial cells, and is essential for both physiological and pathological angiogenesis. Disruption of this gene in mice resulted in abnormal embryonic blood vessel formation. This gene is upregulated in many known tumors and its expression is correlated with tumor stage and progression. Elevated levels of this protein are found in patients with POEMS syndrome, also known as Crow-Fukase syndrome. Allelic variants of this gene have been associated with microvascular complications of diabetes 1 (MVCD1) and atherosclerosis. Alternatively spliced transcript variants encoding different isoforms have been described. There is also evidence for alternative translation initiation from upstream non-AUG (CUG) codons resulting in additional isoforms. A recent study showed that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is antiangiogenic. Expression of some isoforms derived from the AUG start codon is regulated by a small upstream open reading frame, which is located within an internal ribosome entry site. The levels of VEGF are increased during infection with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2),

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

thus promoting inflammation by facilitating recruitment of inflammatory cells, and by increasing the level of angiopoietin II (Ang II), one of two products of the SARS-CoV-2 binding target, angiotensin-converting enzyme 2 (ACE2). In turn, Ang II facilitates the elevation of VEGF, thus forming a vicious cycle in the release of inflammatory cytokines.

Applications	ELISA, CLIA, CG
Notes	For research use only, not for use in diagnostic procedure.
Storage	Short term (up to 7 days) store at 2–8 centigrade. Long term, store at - 20 centigrade. Avoid multiple freeze/thaw cycles.
Storage Buffer	Supplied as a 0.2 µm filtered solution of PBS, pH7.4


GENE INFORMATION

Gene Name	VEGFA vascular endothelial growth factor A [Homo sapiens (human)]
Official Symbol	VEGFA
Synonyms	VEGFA; vascular endothelial growth factor A; vascular endothelial growth factor , VEGF; VEGF A; VPF; vascular permeability factor; VEGF; MVCD1; MGC70609;
Gene ID	7422
mRNA Refseq	NM_001025366
Protein Refseq	NP_001020537
MIM	192240

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID**P15692****SDS-PAGE** Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA