

Active Recombinant Mouse Cxcl2 Protein (73 aa)

Cat. No. Cxcl2-020C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse Cxcl2 Protein without tag was expressed in E. coli.
Species	Mouse
Source	E.coli
ProteinLength	73
Description	Macrophage Inflammatory Protein 2 (MIP-2) was originally identified as a heparin binding protein secreted from a murine macrophage cell line in response to endotoxin stimulation. Based on its protein and DNA sequences, MIP-2 is a member of the alpha (CXC) subfamily of chemokines. Similarly to other alpha chemokines, murine MIP-2 is a potent neutrophil attractant and activator. MIP-2 and KC can bind the murine interleukin 8 type B receptor homologue with high affinity. The expression of MIP-2 was found to be associated with neutrophil influx in pulmonary inflammation and glomerulonephritis, suggesting that MIP-2 may contribute to the pathogenesis of inflammatory diseases.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	Fully biologically active when compared to standard. Determined by its ability to chemoattract total human neutrophils using a concentration range of 1.0-10.0 ng/mL, corresponding to a Specific Activity of 1×10^5 IU/mg.
Molecular Mass	7.8 kDa, a single, non-glycosylated polypeptide chain containing 73 amino acids.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	AVVASELRCQCLKTLPRVDFKNIQSLSVTPPGPHCAQTEVIATLKGGQKVCLDPEAP LVQKIIQKILNKGKAN
Endotoxin	Less than 1 EU/mg of rMuMIP-2/CXCL2 as determined by LAL method.
Purity	>97% by SDS-PAGE and HPLC analyses.
Storage	This lyophilized preparation is stable at 2-8 centigrade, but should be kept at -20 centigrade for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 centigrade. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 to -70 centigrade. Avoid repeated freeze/thaw cycles.
Storage Buffer	Lyophilized from a 0.2mm filtered concentrated solution in 20mM PB, pH 7.4, 150mM NaCl.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	Cxcl2 chemokine (C-X-C motif) ligand 2 [<i>Mus musculus</i>]
Official Symbol	Cxcl2
Synonyms	CXCL2; chemokine (C-X-C motif) ligand 2; C-X-C motif chemokine 2; macrophage inflammatory protein 2; small inducible cytokine subfamily, member 2; GROb; Gro2; Mip2; Scyb; MIP-2; Scyb2; MIP-2a; Mgsa-b; CINC-2a;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	20310
mRNA Refseq	NM_009140
Protein Refseq	NP_033166
UniProt ID	P10889

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA