

Active Recombinant Human CCL7 Protein (76 aa)

Cat. No. CCL7-069C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human CCL7 Protein without tag was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	76
Description	MCP2 and CCL7 are two monocyte chemotactic proteins produced by human MG63 osteosarcoma cells. Both MCP2 and CCL7 are members of the CC family of chemokines and share 62% and 71% amino acid sequence identity, respectively, with MCP1. CCL7 also shares 58% amino acid identity with MCP2. Similarly to other CC chemokines, all three MCP proteins are monocyte chemoattractants. In addition, the three MCPs can chemoattract activated NK cells as well as CD4+ and CD8+ T lymphocytes. All three cytokines have also been shown to attract eosinophils and induce histamine secretion from basophils.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	Fully biologically active when compared to standard. Determined by its ability to chemoattract human peripheral blood monocytes using a concentration range of 10.0-100.0 ng/mL, corresponding to a Specific Activity of 1×10^4 IU/mg.
Molecular Mass	9.0 kDa, a single, non-glycosylated polypeptide chain containing 76 amino acids.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	QPVGINSTTCCYRFINKKIPKQRLESYRRTTSSHCPREAVIFKTKLDKEICADPTQK WVQDFMKHLDKKTQTPKL
Endotoxin	Less than 1 EU/mg of rHuMCP-3/CCL7 as determined by LAL method.
Purity	>97% by SDS-PAGE and HPLC analyses.
Storage	This lyophilized preparation is stable for several weeks at 2-8 centigrade, but should be kept at -20 centigrade for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 centigrade. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 to -70 centigrade. Avoid repeated freeze/thaw cycles.
Storage Buffer	Lyophilized from a 0.2mm filtered concentrated solution in 20mM PB, pH 7.4, 150mM NaCl.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/ml. Stock solutions should be apportioned into working aliquots and stored at < -20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	CCL7 chemokine (C-C motif) ligand 7 [Homo sapiens]
Official Symbol	CCL7
Synonyms	CCL7; chemokine (C-C motif) ligand 7; SCYA6, SCYA7, small inducible cytokine A7 (monocyte chemotactic protein 3); C-C motif chemokine 7; FIC; MARC; MCP 3; MCP3; monocyte chemoattractant protein 3; monocyte chemotactic protein 3; NC28;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

small-inducible cytokine A7; small inducible cytokine A7 (monocyte chemotactic protein 3); MCP-3; SCYA6; SCYA7; MGC138463; MGC138465;

Gene ID	6354
mRNA Refseq	NM_006273
Protein Refseq	NP_006264
MIM	158106
UniProt ID	P80098

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA