

Active Recombinant Human CXCL8 Protein (72 aa)

Cat. No. CXCL8-108C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human CXCL8 Protein without tag was expressed in E. coli.
Species	Human
Source	E.coli
ProteinLength	72
Description	Interleukin 8/CXCL8 was originally discovered and purified independently by a number of laboratories as a neutrophil chemotactic and activating factor. It was also referred to as neutrophil chemotactic factor (NCF), neutrophil activating protein (NAP), monocytederived neutrophil chemotactic factor (MDNCF), T-lymphocyte chemotactic factor (TCF), granulocyte chemotactic protein (GCP) and leukocyte adhesion inhibitor (LAI). Many cell types, including monocyte/macrophages, T cells, neutrophils, fibroblasts, endothelial cells, keratinocytes, hepatocytes, chondrocytes, and various tumor cell lines, can produce CXCL8 in response to a wide variety of pro-inflammatory stimuli such as exposure to IL-1, TNF, LPS, and viruses. CXCL8 is a member of the alpha (C-X-C) subfamily of chemokines, which also includes platelet factor 4, GRO, IP-10, etc.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	Fully biologically active when compared to standard. Determined by its ability to chemoattract human peripheral blood neutrophils using a concentration range of 10.0-100.0 ng/mL, corresponding to a Specific Activity of 1×10^4 IU/mg.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	8,453 Da, a single, non-glycosylated polypeptide chain containing 72 amino acids.
AA Sequence	SAKELRCQCIKTYSKPFHPKFIKELRVIESGPHCANTEIIVKLSDGRELCCLDPKENWV QRVVEKFLKRAENS
Endotoxin	Less than 1 EU/mg of rHuIL-8 as determined by LAL method.
Purity	>95% by SDS-PAGE and HPLC analyses.
Storage	This lyophilized preparation is stable at 2-8 centigrade, but should be kept at -20 centigrade for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 centigrade. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 to -70 centigrade. Avoid repeated freeze/thaw cycles.
Storage Buffer	Lyophilized from a 0.2mm filtered concentrated solution in PBS, pH 7.4.
Reconstitution	We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20 centigrade. Further dilutions should be made in appropriate buffered solutions.

GENE INFORMATION

Gene Name	CXCL8 C-X-C motif chemokine ligand 8 [Homo sapiens (human)]
Official Symbol	CXCL8
Synonyms	CXCL8; C-X-C motif chemokine ligand 8; IL8; NAF; GCP1; LECT; LUCT; NAP1; GCP-1; LY NAP; MDNCF; MONAP; NAP-1; SCYB8; interleukin-8; T-cell chemotactic

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

factor; alveolar macrophage chemotactic factor I; beta endothelial cell-derived neutrophil activating peptide; beta-thromboglobulin-like protein; chemokine (C-X-C motif) ligand 8; emoctakin; granulocyte chemotactic protein 1; interleukin 8; lung giant cell carcinoma-derived chemotactic protein; lymphocyte derived neutrophil activating peptide; monocyte-derived neutrophil chemotactic factor; monocyte-derived neutrophil-activating peptide; neutrophil-activating peptide 1; small inducible cytokine subfamily B, member 8; tumor necrosis factor-induced gene 1

Gene ID	3576
mRNA Refseq	NM_000584
Protein Refseq	NP_000575
MIM	146930
UniProt ID	P10145