

Active Recombinant Rat Ccl20 Protein (71 aa)

Cat. No. Ccl20-170C Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant rat MIP-3 α /CCL20 produced in HEK293 cells is a polypeptide chain containing 71 amino acids. A fully biologically active molecule, rrMIP-3 α /CCL20 has a molecular mass of 8.2 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques.
Species	Rat
Source	HEK293
ProteinLength	71
Description	Macrophage Inflammatory Protein-3 (MIP-3 α), also known as chemokine (C-C motif) ligand 20 (CCL20) or liver activation regulated chemokine (LARC), is a small cytokine belonging to the CC chemokine family. It is strongly chemotactic for lymphocytes and weakly attracts neutrophils. MIP-3 α is implicated in the formation and function of mucosal lymphoid tissues via chemoattraction of lymphocytes and dendritic cells toward the epithelial cells surrounding these tissues. MIP-3 α is expressed in the liver, lymph nodes, appendix, PBL and lung and can signal through the CCR6 receptor.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	The EC ₅₀ value of rat MIP-3 α /CCL20 on Ca ²⁺ mobilization assay in CHO-K1/G α 15/rCCR6 cells (human G α 15 and rat CCR6 stably expressed in CHO-K1 cells) is less than 0.6 μ g/mL.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	8.2 kDa, observed by reducing SDS-PAGE.
AA Sequence	ASNFDCCCLTYTKNVYHHARNFVGFTTQMADEACDINAIIIFHLKSKRSVCADPKQIWW KRILHLLSLRTKKM
Endotoxin	< 0.2 EU/μg, determined by LAL method.
Purity	> 98% as analyzed by SDS-PAGE.
Storage	Lyophilized recombinant Rat MIP-3 α/CCL20 remains stable up to 6 months at -80 centigrade from date of receipt. Upon reconstitution, recombinant Rat MIP-3 α/CCL20 should be stable up to 1 week at 4 centigrade or up to 2 months at -20 centigrade.
Storage Buffer	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 μg/mL.

GENE INFORMATION

Gene Name	Ccl20 chemokine (C-C motif) ligand 20 [<i>Rattus norvegicus</i>]
Official Symbol	Ccl20
Synonyms	CCL20; chemokine (C-C motif) ligand 20; C-C motif chemokine 20; MIP-3-alpha; CC chemokine LARC; CC chemokine ST38; beta chemokine exodus-1; beta-chemokine exodus-1; small-inducible cytokine A20; small inducible cytokine subfamily A20; macrophage inflammatory protein 3 alpha; liver and activation-regulated chemokine; ST38; Scya20;
Gene ID	29538

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_019233
Protein Refseq	NP_062106
UniProt ID	P97884

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA