

Active Recombinant Rat Ccl2 Protein (125 aa)

Cat. No. Ccl2-172C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant rat MCP-1/CCL2 produced in HEK293 cells is a polypeptide chain containing 125 amino acids. A fully biologically active molecule, rrMCP-1/CCL2 has a molecular mass of 10-15 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques.
Species	Rat
Source	HEK293
ProteinLength	125
Description	Chemokine (C-C motif) ligand 2 (CCL2) is also referred to as monocyte chemotactic protein 1 (MCP1) and small inducible cytokine A2. CCL2 is a small cytokine that belongs to the CC chemokine family. CCL2 recruits monocytes, memory T cells, and dendritic cells to the sites of inflammation produced by either tissue injury or infection. CCL2 is implicated in the pathogenesises of several diseases characterized by monocytic infiltrates, such as psoriasis, rheumatoid arthritis and atherosclerosis. CCL2 is anchored in the plasma membrane of endothelial cells by glycosaminoglycan side chains of proteoglycans. CCL2 is primarily secreted by monocytes, macrophages and dendritic cells. CCL2 can signal through the CCR2 receptor.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	The EC50 value of rat MCP-1/CCL2 on Ca ²⁺ mobilization assay in CHO-K1/G

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	α 15/rCCR2 cells (human G α 15 and rat CCR2 stably expressed in CHO-K1 cells) is less than 0.3 μ g/mL.
Molecular Mass	10~15 kDa, observed by reducing SDS-PAGE.
AA Sequence	QPDVNAPLTCCYSFTGKMIPMSRLENYKRITSSRCPKEAVVFVTKLKREICADPNK EWVQKYIRKLDQNQVRSETTVFYKIASTLRTSAPLNVNLTHKSEANASTLFSTTTSSST SVEVTSMTEN
Endotoxin	< 0.2 EU/ μ g, determined by LAL method.
Purity	> 98% as analyzed by SDS-PAGE.
Storage	Lyophilized recombinant Rat MCP-1/CCL2 remains stable up to 6 months at -80 centigrade from date of receipt. Upon reconstitution, Rat MCP-1/CCL2 should be stable up to 1 week at 4 centigrade or up to 3 months at -20 centigrade.
Storage Buffer	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 μ g/mL.

GENE INFORMATION

Gene Name	Ccl2 chemokine (C-C motif) ligand 2 [<i>Rattus norvegicus</i>]
Official Symbol	Ccl2
Synonyms	CCL2; chemokine (C-C motif) ligand 2; C-C motif chemokine 2; small inducible gene JE; small inducible cytokine A2; small-inducible cytokine A2; monocyte chemotactic protein 1; monocyte chemoattractant protein 1; monocyte chemoattractant protein-1; immediate-early serum-responsive JE protein; immediate-early serum-responsive

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

protein JE; MCP-1; Scya2; Sigje;

Gene ID 24770**mRNA Refseq** NM_031530**Protein Refseq** NP_113718**UniProt ID** P14844 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA