

Active Recombinant Human CCL23 Protein (75 aa)

Cat. No. CCL23-217C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human MPIF-1/CCL23 (aa46-120) produced in CHO cells is a single polypeptide chain containing 75 amino acids. A fully biologically active molecule, rhMPIF-1/CCL23 (aa46-120) has a molecular mass of 11.3 kDa analyzed by reducing SDS-PAGE and is obtained by chromatographic techniques.
Species	Human
Source	CHO
ProteinLength	75
Description	Myeloid progenitor inhibitory factor 1 (MPIF-1), also known as Chemokine (C-C motif) ligand 23 (CCL23) is a small cytokine belonging to the CC chemokine family. MPIF-1 is predominantly expressed in lung and liver tissue, but is also found in bone marrow and placenta. It is also expressed in some cell lines of myeloid origin. It is highly chemotactic for resting T cells and monocytes and slightly chemotactic for neutrophils. MPIF-1 has been shown to inhibit colony formation of bone marrow myeloid immature progenitors. It has also been attributed to an inhibitory activity on hematopoietic progenitor cells. MPIF-1 is a ligand for the chemokine receptor CCR1.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	The EC ₅₀ value of human MPIF-1/CCL23 (aa46-120) on Ca ²⁺ mobilization assay in CHO-K1/Gα15/hCCR1 cells (human Gα15 and human CCR1 stably expressed in CHO-K1 cells) is less than 1

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	µg/mL.
Molecular Mass	11.3 kDa, observed by reducing SDS-PAGE.
AA Sequence	RFHATSADCCISYTPRSIPCSLLESYFETNSECSKPGVIFLTKKGRRFCANPSDKQVQVCVRMLKLDTRIKTRKN
Endotoxin	< 0.2 EU/µg, determined by LAL method.
Purity	> 95% as analyzed by SDS-PAGE.
Storage	Lyophilized recombinant Human MPIF/CCL23 (aa46-120) remains stable up to 6 months at -80 centigrade from date of receipt. Upon reconstitution, Human MPIF/CCL23 (aa46-120) should be stable up to 1 week at 4 centigrade or up to 3 months at -20 centigrade.
Storage Buffer	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O or PBS at 100 µg/mL.

GENE INFORMATION

Gene Name	CCL23 chemokine (C-C motif) ligand 23 [Homo sapiens]
Official Symbol	CCL23
Synonyms	CCL23; chemokine (C-C motif) ligand 23; SCYA23, small inducible cytokine subfamily A (Cys Cys), member 23; C-C motif chemokine 23; Ckb 8; CKb8; MIP 3; MPIF 1; hmrb-2a; C6 beta-chemokine; small-inducible cytokine A23; macrophage inflammatory protein 3; myeloid progenitor inhibitory factor 1; small inducible cytokine subfamily A (Cys-Cys), member 23; MIP3; Ckb-8; MIP-3; MPIF-1; SCYA23; Ckb-8-1;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

CK-BETA-8;

Gene ID

6368

mRNA Refseq

NM_005064

Protein Refseq

NP_005055

MIM

602494

UniProt ID

P55773

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA