

Active Recombinant Human TNFSF13B Protein

Cat. No. TNFSF13B-299T **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human TNFSF13B Protein without tag was expressed in CHO.
Species	Human
Source	CHO
Description	B-cell activating factor, also known as BAFF, TALL-1, TNAK, and zTNF4, is a member of the TNF ligand superfamily designated TNFSF13B. Produced by macrophages, dendritic cells, and T lymphocytes, BAFF promotes the survival of B cells and is essential for B cell maturation. BAFF binds to three TNF receptor superfamily members: B-cell maturation antigen (BCMA/TNFRSF17), transmembrane activator and calcium-modulator and cyclophilin ligand interactor (TACI/TNFRSF13B) and BAFF receptor (BAFF R/BR3/TNFRSF 13C). These receptors are type III transmembrane proteins lacking a signal peptide. Whereas TACI and BCMA bind BAFF and another TNF superfamily ligand, APRIL (a proliferation-inducing ligand), BAFF R selectively binds BAFF. The BAFF R extracellular domain lacks the TNF receptor canonical cysteine-rich domain (CRD) and contains only a partial CRD with four cysteine residues. Human and mouse BAFF R share 56% aa sequence identity. BAFF R is highly expressed in spleen, lymph node and resting B cells. It is also expressed at lower levels in activated B cell, resting CD4+ T cells, thymus and peripheral blood leukocytes.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Bio-activity	ED50< 20 ng/mL, determined by dose-dependent mitogenic activity on human RPMI 8226 cells, corresponding to a specific activity of >5.0 4 units/mg.
Molecular Mass	17kDa, observed by non-reducing SDS-PAGE.
AA Sequence	AVQGPEETVTQDCLQLIADSETPTIQKGSYTFVPWLLSFKRGSAALEEKENKILVKETG YFFIYGQVLYTDKTYAMGHLIQRKKVHVFGDELSTLFRICIQNMPETLPNNSCYSA GIAKLEEGDELQLAIPRENAQISLDGDVTFFGALKLL
Endotoxin	< 0.2 EU/μg, determined by LAL method.
Purity	> 95% as analyzed by SDS-PAGE and HPLC.
Storage	Lyophilized recombinant human B-Cell Activating Factor (BAFF) remains stable up to 6 months at -80 centigrade from date of receipt. Upon reconstitution, rh-BAFF should be stable up to 1 week at 4 centigrade or up to 2 months at -20 centigrade.
Storage Buffer	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH2O or PBS at 100 μg/mL.
GENE INFORMATION	
Gene Name	TNFSF13B tumor necrosis factor (ligand) superfamily, member 13b [Homo sapiens]
Official Symbol	TNFSF13B
Synonyms	TNFSF13B; tumor necrosis factor (ligand) superfamily, member 13b; TNFSF20; tumor necrosis factor ligand superfamily member 13B; BAFF; BLYS; CD257; TALL 1; TALL1; THANK; delta BAFF; Delta4 BAFF; B-lymphocyte stimulator; B-cell-activating factor; ApoL related ligand TALL-1; TNF homolog that activates apoptosis; dendritic

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

cell-derived TNF-like molecule; tumor necrosis factor-like protein ZTNF4; TNF and ApoL-related leukocyte expressed ligand 1; tumor necrosis factor (ligand) superfamily, member 20; DTL; ZTNF4; TALL-1;

Gene ID [10673](#)

mRNA Refseq [NM_001145645](#)

Protein Refseq [NP_001139117](#)

MIM [603969](#)

UniProt ID [Q9Y275](#)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA