

Active Recombinant Human CXCL8 Protein (8-79aa, 72 aa)

Cat. No. CXCL8-349C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human I Interleukin-8/CXCL8 (rhIL-8) produced in E. coli is a single non-glycosylated polypeptide chain containing 72 amino acids. A fully biologically active molecule, rhIL-8 has a molecular mass of 8.4 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques.
Species	Human
Source	E.coli
ProteinLength	72
Description	Interleukin-8 is one of the first discovered chemokines, and belongs to the CXCL family, in which the first two conserved cysteines are separated by one residue. In vivo, IL-8 exists in two forms: 77 a.a. produced by endothelial cells, and the more active 72 a.a. produced by monocytes. The receptors of IL-8 are the seven-helical G-protein coupled receptors CXCR1 and CXCR2, exclusively expressed on neutrophils. The functions of IL-8 are to induce rapid changes in cellular shape, activate the integrins, and release the granule contents of neutrophils. Thus, IL-8 can enhance the antimicrobial actions of defense cells.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	ED50 < 20 ng/mL, measured by the FLIPR assay using CHO cells transfected with human CXCR1, the receptor of human CXCL8, corresponding to a specific activity of > 5 × 10 ⁴ units/mg.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	8.4 kDa, observed by reducing SDS-PAGE.
AA Sequence	SAKELRCQCIKTYSKPFHPKFIKELRVIESGPHCANTEIIVKLSDGRELCCLDPKENWV QRVVEKFLKRAENS
Endotoxin	< 0.2 EU/μg, determined by LAL method.
Purity	> 95% as analyzed by SDS-PAGE and HPLC.
Storage	Lyophilized recombinant human Interleukin-8/CXCL8 (rhIL-8) remains stable up to 6 months at -80 centigrade from date of receipt. Upon reconstitution, rhIL-8 remains stable up to 2 weeks at 4 centigrade or up to 3 months at -20 centigrade.
Storage Buffer	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O at 100 μg/mL.

GENE INFORMATION

Gene Name	CXCL8 C-X-C motif chemokine ligand 8 [Homo sapiens (human)]
Official Symbol	CXCL8
Synonyms	CXCL8; C-X-C motif chemokine ligand 8; IL8; NAF; GCP1; LECT; LUCT; NAP1; GCP-1; LYNAP; MDNCF; MONAP; NAP-1; SCYB8; interleukin-8; T-cell chemotactic factor; alveolar macrophage chemotactic factor I; beta endothelial cell-derived neutrophil activating peptide; beta-thromboglobulin-like protein; chemokine (C-X-C motif) ligand 8; emoctakin; granulocyte chemotactic protein 1; interleukin 8; lung giant cell carcinoma-derived chemotactic protein; lymphocyte derived neutrophil activating peptide; monocyte-derived neutrophil chemotactic factor; monocyte-derived neutrophil-activating peptide; neutrophil-activating peptide 1; small inducible cytokine

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

subfamily B, member 8; tumor necrosis factor-induced gene 1

Gene ID 3576**mRNA Refseq** NM_000584**Protein Refseq** NP_000575**MIM** 146930**UniProt ID** P10145 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA