

Active Recombinant Human FGF2 Protein (146 aa)

Cat. No. FGF2-401F **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human Fibroblast Growth Factor-basic (146 a.a.) (rhFGF-basic) produced in E. coli is a single non-glycosylated polypeptide chain containing 146 amino acids. A fully biologically active molecule, rhFGF-basic has a molecular mass of 16.4 kDa analyzed by reducing SDS-PAGE and is obtained by proprietary chromatographic techniques.
Species	Human
Source	E.coli
ProteinLength	146
Description	Fibroblast Growth Factor-basic (FGF-basic), also known as FGF-2, is a pleiotropic cytokine and one of the prototypic members of the heparin-binding FGF family. Like other FGF family members, FGF-basic has the β trefoil structure. In vivo, FGF-basic is produced by a variety of cells, including cardiomyocytes, fibroblasts, and vascular cells. FGF-basic regulates a variety of processes including cell proliferation, differentiation, survival, adhesion, motility, apoptosis, limb formation and wound healing. FGF-basic can be tumorigenic due to its role in angiogenesis and blood vessel remodeling. The angiogenic effects of FGF-basic can produce beneficial cardioprotection during acute heart injury.
Form	Sterile Filtered White lyophilized (freeze-dried) powder.
Bio-activity	ED50 < 0.25 ng/mL, measured by the cell proliferation assay using 3T3 cells,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

corresponding to a specific activity of $> 4 \times 10^6$ units/mg.

Molecular Mass	16.4 kDa, observed by reducing SDS-PAGE.
AA Sequence	PALPEDGGSGAFPPGHFKDPKRLYCKNGGFFLRIHPDGRVDGVREKSDPHIKLQLQ AEERGVSISIKGVCANRYLAMKEDGRLLASKCVTDECFFFERLESNNYNTYRSRKYT SWYVALKRTGQYKLGSKTGPQKAILFLPMSAKS
Endotoxin	< 0.2 EU/μg, determined by LAL method.
Purity	> 95% by SDS-PAGE analysis.
Storage	Lyophilized recombinant human Fibroblast Growth Factor-basic (146 a.a.) (rhFGF-basic) remains stable up to 6 months at -80 centigrade from date of receipt. Upon reconstitution, rhFGF-basic remains stable up to 2 weeks at 4 centigrade or up to 3 months at -20 centigrade.
Storage Buffer	Lyophilized after extensive dialysis against PBS.
Reconstitution	Reconstituted in ddH ₂ O at 50 μg/mL.

GENE INFORMATION

Gene Name	FGF2 fibroblast growth factor 2 (basic) [Homo sapiens]
Official Symbol	FGF2
Synonyms	FGF2; fibroblast growth factor 2 (basic); FGFB; fibroblast growth factor 2; prostatropin; heparin-binding growth factor 2; basic fibroblast growth factor bFGF; BFGF; FGF-2; HBGF-2;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	2247
mRNA Refseq	NM_002006
Protein Refseq	NP_001997
MIM	134920
UniProt ID	P09038

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA