

Active Recombinant Human IL7 Protein (Asp26-His177), GMP Grade, Animal-Free

Cat. No. IL7-23HG Lot. No. (See product label)

SPECIFICATION

Product Overview	GMP recombinant human IL-7 protein (Asp26-His177) with an N-terminal Met was expressed in E. coli, produced in an Animal-Free laboratory and manufactured and tested under cGMP guidelines.
Species	Human
Source	E.coli
ProteinLength	Asp26-His177
Description	The protein encoded by this gene is a cytokine important for B and T cell development. This cytokine and the hepatocyte growth factor (HGF) form a heterodimer that functions as a pre-pro-B cell growth-stimulating factor. IL7 is found to be a cofactor for V(D)J rearrangement of the T cell receptor beta (TCRB) during early T cell development. This cytokine can be produced locally by intestinal epithelial and epithelial goblet cells, and may serve as a regulatory factor for intestinal mucosal lymphocytes. IL7 plays an essential role in lymphoid cell survival, and in the maintenance of naive and memory T cells. Alternative splicing results in multiple transcript variants encoding distinct isoforms. Additional splice variants have been described but their presence in normal tissues has not been confirmed. Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection can be a potent inducer of proinflammatory cytokines and chemokines which may defend against the infection, but may also mediate destructive lung injury. Elevated serum IL7 levels,

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

	together with several other circulating cytokines and chemokines, has been found to be associated with the severity of Coronavirus Disease 19 (COVID-19).
Bio-activity	Measured in a cell proliferation assay using PHA-activated human peripheral blood lymphocytes (PBL). The ED50 for this effect is 0.100-0.500 ng/mL. The specific activity of Recombinant Human IL-7 is >1.00 x 10 ⁸ units/mg, which is calibrated against the human IL-7 reference standard.
Molecular Mass	Predicted Molecular Mass: 17 kDa SDS-PAGE: 17 kDa, under reducing conditions
N-terminal Sequence Analysis	Met-Asp26-(Cys)-Asp-Ile-Glu-Gly-Lys-Asp-Gly
Endotoxin	<0.10 EU/ µg of the protein by the LAL method.
Purity	>97%, by SDS-PAGE visualized with Silver Staining and quantitative densitometry by Coomassie® Blue Staining.
Storage	Use a manual defrost freezer and avoid repeated freeze-thaw cycles. A minimum of 12 months when stored at ≤ -20 centigrade as supplied. Refer to lot specific COA for the Use by Date. 1 month, 2 to 8 centigrade under sterile conditions after reconstitution. 3 months, ≤ -20 centigrade under sterile conditions after reconstitution.
Storage Buffer	Lyophilized from a 0.2 µm filtered solution in PBS with Trehalose.
Reconstitution	Reconstitute at 100-500 µg/mL in PBS.
Shipping	The product is shipped with polar packs.

GENE INFORMATION

Gene Name	IL7 interleukin 7 [Homo sapiens]
Official Symbol	IL7
Synonyms	IL7; interleukin 7; interleukin-7; IL 7; IL-7;
Gene ID	3574
mRNA Refseq	NM_000880
Protein Refseq	NP_000871
MIM	146660
UniProt ID	P13232

Bioactivity




Tel: 1-631-559-9269 1-516-512-3133


Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA

SDS-PAGE



Mass Spectrometry



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA