

Recombinant Mouse Interleukin 25

Cat. No. IL25-199M Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Mouse IL17E produced in <i>E. Coli</i> is a homodimeric, non-glycosylated polypeptide chain containing a total of 306 amino acids and having a molecular mass of 34.9 kDa. The Mouse IL-17E is purified by proprietary chromatographic techniques.
-------------------------	--

Species	Mouse
----------------	-------

Source	E.coli
---------------	--------

Description

IL-25 also called IL-17E cytokine has a sequence similarity with IL17. IL-17E induces NF-kappaB activation, and stimulates the production of IL-8. IL17E and IL17B are ligands for the cytokine receptor IL17BR. IL-25 is a proinflammatory cytokine favoring Th2-type immune response. The upregulation of costimulation-induced IL-17E receptors and release of cytokines and chemokines from IL-17E treated costimulated Th cells are differentially regulated by intracellular JNK, p38 MAPK and NF-kappaB activity. Blocking interleukin-25 prevents airway hyperresponsiveness, a critical feature of clinical asthma. IL25 produced by innate effector eosinophils and basophils increase the allergic inflammation by enhancing the maintenance and functions of TSLP-DC activated adaptive Th2 memory cells. Over expression of IL-25 up-regulates gene expression of Th2 cytokines and induces growth retardation, jaundice, and multiorgan inflammation in a transgenic mouse model. IL-25 contributes to the induction and maintenance of eosinophilic inflammation by acting on lung fibroblasts which supports the fact that IL-17E is an important factor in asthma pathophysiology. IL-17E operates by amplifying TH2 cell-mediated allergic airway inflammation but doesn't induce allergic inflammation in vivo.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Amino Acid Sequence	VSLRIQEGCSHLPSCCPSKEQEPPEEWLKWSSASVSPPEPLSHTTHAESCRAKDGPLNSRAIS PWSYELDRDLNRVPQDLYHARCLCPHCVSLQTGSHM DPLGNSVPLY H NQTV FY RRPCHGEEGTHRRYCLERRLYRVSLACVCVRP RVMA.
Purity	Greater than 95.0% as determined by: (a) Analysis by RP-HPLC. (b) Analysis by SDS-PAGE.
Formulation	IL17E was lyophilized from a concentrated (1mg/ml) solution containing no additives.
Solubility	It is recommended to reconstitute the lyophilized Mouse IL17E in sterile 18MΩ-cm H2O not less than 100g/ml, which can then be further diluted to other aqueous solutions.
Stability	Lyophilized Murine IL17E although stable at room temperature for 3 weeks, should be stored desiccated below -18°C. Upon reconstitution IL17E should be stored at 4°C between 2-7 days and for future use below -18°C. For long term storage it is recommended to add a carrier protein (0.1% HSA or BSA). Please prevent freeze-thaw cycles.

GENE INFORMATION

Gene Name	IL25 interleukin 25 [<i>Mus musculus</i>]
Synonyms	IL25; interleukin 25; IL17e; IL-17E
Gene ID	140806
mRNA Refseq	NM_080729
Protein Refseq	NP_542767

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

UniProt ID	Q8VHC9
Chromosome Location	14 C3
Pathway	Cytokine-cytokine receptor interaction
Function	cytokine activity;interleukin-17E receptor binding

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA