

Recombinant Human proteasome (prosome, macropain) subunit, beta type, 8, His-tagged

Cat. No. PSMB8-391H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	PSMB8, 73-276aa, Human, His tag, E.coli?
Species	Human
Source	E.coli
ProteinLength	73-276 a.a.
Description	PSMB8, also known as LMP7, belongs to the peptidase T1B family. The proteasome is a multicatalytic proteinase complex with a highly ordered ring-shaped 20S core structure. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMB8 is located in the class II region of the MHC (major histocompatibility complex). Replacement of PSMB5 by PSMB8 increases the capacity of the immunoproteasome to cleave model peptides after hydrophobic and basic residues. This protein acts as a major component of interferon gamma-induced sensitivity. It plays a key role in apoptosis via the degradation of the apoptotic inhibitor MCL1. Recombinant human PSMB8 protein, fused to His-tag at N-terminus, was expressed in E.coli.
Form	Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol
Molecular Mass	25.4 kDa (229aa)

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	MGSSHHHHHH SSGLVPRGSH MGSMTTTLA FKFQHGVIAA VDSRASAGSY ISALRVNKVI EINPYLLGTM SGCAADCQYW ERLLAKECRL YYLRNGERIS VSAASKLLSN MMCQYRGMGL SMGSMICGWD KKGPGLYYVD EHGTRLSGNM FSTGSGNTYA YGVMDSGYRP NLSPEEAYDL GRRAIAYATH RDSYSGGVVN MYHMKEDGWV KVESTDVSDL LHQYREANQ
Purity	>85% by SDS - PAGE
Storage	Can be stored at +4C short term (1-2 weeks). For long term storage, aliquot and store at -20C or -70C. Avoid repeated freezing and thawing cycles.
Concentration	0.5 mg/ml (determined by Bradford assay)
GENE INFORMATION	
Gene Name	PSMB8 proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional peptidase 7) [Homo sapiens]
Official Symbol	PSMB8
Synonyms	PSMB8; proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional peptidase 7); LMP7, proteasome (prosome, macropain) subunit, beta type, 8 (large multifunctional protease 7); proteasome subunit beta type-8; beta5i; D6S216E; PSMB5i; RING10; macropain subunit C13; proteasome subunit Y2; protease component C13; proteasome component C13; proteasome-related gene 7; proteasome subunit beta 5i; low molecular mass protein 7; low molecular weight protein 7; proteasome catalytic subunit 3i; really interesting new gene 10 protein; multicatalytic endopeptidase complex subunit C13; JMP; ALDD; LMP7; NKJO; D6S216; MGC1491;
Gene ID	5696

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_004159
Protein Refseq	NP_004150
MIM	177046
UniProt ID	P28062
Chromosome Location	6p21.3
Pathway	APC/C-mediated degradation of cell cycle proteins, organism-specific biosystem; APC/C:Cdc20 mediated degradation of Securin, organism-specific biosystem; APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; APC/C:Cdh1 mediated degradation of Cdc20 and other APC/C:Cdh1 targeted proteins in late mitosis/early G1, organism-specific biosystem; Activation of APC/C and APC/C:Cdc20 mediated degradation of mitotic proteins, organism-specific biosystem; Activation of NF-kappaB in B Cells, organism-specific biosystem; Adaptive Immune System, organism-specific biosystem;
Function	peptidase activity; threonine-type endopeptidase activity;