

Recombinant Mouse chemokine (C-C motif) ligand 2, His-tagged

Cat. No. CCL2-559M **Lot. No.** (See product label)

SPECIFICATION

Product Overview	CCL2, 24-148aa, Mouse His tag, E.coli
Species	Mouse
Source	E.coli
ProteinLength	24-148 a.a.
Description	Chemokine (C-C motif) ligand 2 (CCL2) is a small cytokine belonging to the CC chemokine family that is also known as monocyte chemotactic protein-1 (MCP-1). CCL2 recruits monocytes, memory T cells, and dendritic cells to sites of tissue injury and infection. It has been implicated in the pathogenesis of diseases characterized by monocytic infiltrates, like psoriasis, rheumatoid arthritis and atherosclerosis. It binds to chemokine receptors CCR2 and CCR4. Recombinant mouse CCL2 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.
Form	Liquid. In PBS (pH 7.4) containing 10% glycerol.
Molecular Mass	16 kDa (146aa) confirmed by MALDI-TOF
AA Sequence	MGSSHHHHHH SSGLVPRGSH MQPDAVNAPL TCCYSFTSKM IPMSRLESYK RITSSRCPKE AVVFVTKLKR EVCADPKKEW VQTYIKNLDR NQMRSEPTTL FKTASALRSS APLNVKLTRK SEANASTTFS TTSSTSVGV TSVTVN

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	>90% by SDS - PAGE
Storage	Can be stored at +4C short term (1-2 weeks). For long term storage, aliquot and store at -20C or -70C. Avoid repeated freezing and thawing cycles.
Concentration	0.25 mg/ml (determined by Bradford assay)
GENE INFORMATION	
Gene Name	Ccl2 chemokine (C-C motif) ligand 2 [<i>Mus musculus</i>]
Official Symbol	CCL2
Synonyms	CCL2; chemokine (C-C motif) ligand 2; C-C motif chemokine 2; small inducible cytokine A2; small-inducible cytokine A2; monocyte chemotactic protein 1; monocyte chemoattractant protein 1; monocyte chemoattractant protein-1; platelet-derived growth factor-inducible protein JE; JE; HC11; MCAF; MCP1; MCP-1; Scya2; Sigje; SMC-CF; AI323594;
Gene ID	20296
mRNA Refseq	NM_011333
Protein Refseq	NP_035463
Pathway	Chemokine signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem;
Function	CCR2 chemokine receptor binding; CCR2 chemokine receptor binding; G-protein coupled receptor binding; chemokine activity; cytokine activity; heparin binding; protein binding;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA