

Recombinant Canine ACVR1, His-tagged

Cat. No. ACVR1-161C **Lot. No.** (See product label)

SPECIFICATION

Product Overview	A DNA sequence encoding the canine ACVR1 (XP_549615.2) (Met1-Glu123) was expressed with a C-terminal polyhistidine tag.
Species	Canine
Source	Human Cells
ProteinLength	Met1-Glu123
Description	Cytokines and hormones belonging to the transforming growth factor beta (TGFβ) superfamily such as the activin, anti-müllerian hormone (AMH), bone morphogenetic proteins (BMPs), and Nodal, are involved in a variety of physiological processes including growth, cell differentiation, homeostasis, osteogenesis, and apoptosis. They initiate signaling through heteromeric complexes of a type I and a type II serine/threonine kinase receptor. The two type receptors are type I transmembrane proteins and are distinguished by the presence of a glycine/serine-rich juxta-membrane domain found in the type I receptors. Type II receptors are required for binding ligands and recruitment of type I receptors which then form stable complexes, and type I receptors are essential for signaling transduction. ALK2, also termed as ACVR1, ActRI, SKR1, was initially identified as an activin type I receptor because of its ability to bind activin in concert with ActRII or IIB. In addition, ALK2 is also identified as a BMP type I receptor. ACVR1 has been demonstrated to form complex with either the BMP-2/7- bound BMPR-II or ACVR2A/ACVR2B and transduce signals to downstream components R-SMADs. Recombinant soluble ALK2 does not bind

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	activin. Mutations in this gene are associated with fibrodysplasia ossificans progressive.
Form	Lyophilized from sterile PBS, pH7.4.
Molecular Mass	The recombinant canine ACVR1 comprises 114 amino acids and has a predicted molecular mass of 13 kDa. The apparent molecular mass of the protein is approximately 19 kDa in SDS-PAGE under reducing conditions due to glycosylation.
Endotoxin	< 1.0 eu per µg of the protein as determined by the LAL method.
Purity	>95 % as determined by SDS-PAGE
Stability	Samples are stable for up to twelve months from date of receipt at -70°C
Storage	Store it under sterile conditions at -70°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.
Reconstitution	Hardcopy of COA with reconstitution instruction is sent along with the products.

GENE INFORMATION

Gene Name	ACVR1 activin A receptor, type I [Canis lupus familiaris]
Official Symbol	ACVR1
Synonyms	ACVR1; activin A receptor, type I; activin receptor type-1;
Gene ID	478757
Pathway	Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

cytokine receptor interaction, conserved biosystem; TGF-beta signaling pathway,
organism-specific biosystem; TGF-beta signaling pathway, conserved biosystem;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA