

Recombinant Human BTLA protein, hFc-tagged

Cat. No. BTLA-233H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant Human BTLA protein(Q7Z6A9-1)(Lys31-Ser150), fused with C-terminal hFc tag, was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	Lys31-Ser150
Tag	C-hFc
Form	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 5% trehalose is added as protectant before lyophilization.
Bio-activity	Immobilized Human HVEM, His Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Human BTLA, hFc Tag with the EC50 of 7.2ng/ml determined by ELISA.
Molecular Mass	The protein has a predicted MW of 40.5 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE; > 95% as determined by HPLC


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127


 45-1 Ramsey Road, Shirley, NY 11967, USA

Storage	Reconstituted protein stable at -80°C for 12 months, 4°C for 1 week. Use a manual defrost freezer and avoid repeated freeze-thaw cycles.
Reconstitution	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
GENE INFORMATION	
Gene Name	BTLA B and T lymphocyte associated [Homo sapiens]
Official Symbol	BTLA
Synonyms	BTLA; B and T lymphocyte associated; B- and T-lymphocyte attenuator; BTLA1; CD272; B and T lymphocyte attenuator; B- and T-lymphocyte-associated protein; FLJ16065; MGC129743;
Gene ID	151888
mRNA Refseq	NM_001085357
Protein Refseq	NP_001078826
MIM	607925
UniProt ID	Q7Z6A9

Tris-Bis PAGE



ELISA Data



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA