

Recombinant Human BTK, His-tagged, Active

Cat. No. BTK-255H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant full-length human BTK was expressed by baculovirus in <i>Sf9</i> insect cell/susing a N-terminal His tag. MW = 78 kDa.
Species	Human
Source	Sf9 Cells
Description	BTK (also known as Bruton tyrosine kinase) plays a crucial role in B-lymphocyte differentiation and activation. BTK interacts with SRC homology 3 domains of FYN, LYN and HCK that are activated upon stimulation of B- and T-cell receptors. Defects in the BTK gene cause Agammaglobulinemia, an X-linked immunodeficiency characterized by failure to produce mature B lymphocyte cells and associated with a failure of Ig heavy chain rearrangement. The unique role of BTK makes it a desirable target for potential anti-cancer, anti-inflammatory and anti-viral agents as well as other treatments.
Sequence	Full-length.
Applications	Kinase Assay, Western Blot.
Storage And Stability	Store product at -70°C. For optimal storage, aliquot target into smaller quantities after centrifugation and store at recommended temperature. For most favorable performance, avoid repeated handling and multiple freeze/thaw cycles.

GENE INFORMATION


 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene Name	BTK Bruton agammaglobulinemia tyrosine kinase [Homo sapiens]
Synonyms	BTK; Bruton agammaglobulinemia tyrosine kinase; AT; ATK; BPK; XLA; IMD1; AGMX1; PSCTK1; MGC126261; MGC126262; OTTHUMP00000023676; OTTHUMP00000063593; Bruton tyrosine kinase; B-cell progenitor kinase; tyrosine-protein kinase BTK; agammaglobulinemia tyrosine kinase; dominant-negative kinase-deficient Brutons tyrosine kinase
Gene ID	695
mRNA Refseq	NM_000061
Protein Refseq	NP_000052
MIM	300300
UniProt ID	Q06187
Chromosome Location	Xq21.33-q22
Pathway	B cell receptor signaling pathway; Fc epsilon RI signaling pathway
Function	ATP binding; identical protein binding; kinase activity; metal ion binding; non-membrane spanning protein tyrosine kinase activity; nucleotide binding; protein binding; protein tyrosine kinase activity; transferase activity; zinc ion binding

PH domain of
Bruton's tyrosine
kinase dimer with
bound lipids. Blue
plane shows
hydrocarbon
boundary of the lipid
bilayer.



Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA