

Recombinant Mouse CXCL10 (IP-10)

Cat. No. CXCL10-71M **Lot. No.** (See product label)

SPECIFICATION

Product Overview

Mouse IP-10/CXCL10 was produced in yeast and therefore does not have endotoxin, is naturally folded, and post-translationally modified.

Species

Mouse

Source

Yeast

Description

C-X-C motif chemokine 10 (CXCL10) also known as Interferon gamma-induced protein 10 (IP-10) or small-inducible cytokine B10 is a member of the C-X-C chemokine family. CXCL10 (IP-10) is secreted by several cell types in response to IFN-gamma. These cell types include monocytes, endothelial cells and fibroblasts. CXCL10 (IP-10) has been attributed to several roles, such as chemoattraction for monocytes/macrophages, T cells, NK cells, and dendritic cells, promotion of T cell adhesion to endothelial cells, antitumor activity, and inhibition of bone marrow colony formation and angiogenesis. There have been 17 different C-X-C chemokines described in mammals, that are subdivided into two categories, those with a specific amino acid sequence (or motif) of glutamic acid-leucine-arginine (or ELR for short) immediately before the first cysteine of the C-X-C motif (ELR-positive), and those without an ELR motif (ELR-negative). ELR-positive C-X-C chemokines specifically induce the migration of neutrophils, and interact with chemokine receptors CXCR1 and CXCR2. C-X-C chemokines that lack the ELR motif are chemoattractant for lymphocytes. CXCL10 (IP-10) elicits its effects by binding to the cell surface chemokine receptor CXCR3.

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Form	Lyophilized
Molecular Mass	8.7 kDa
AA Sequence	IPLARTVRCN CIHIDGDPVR MRAIGKLEII PASLSCPRVE IIATMKKNDE QRCLNPESKT IKNLMKAFSQ KRSKRAP (77)
Applications	The mouse IP-10/CXCL10 protein can be used in cell culture, as an IP-10/CXCL10 ELISA Standard, and as a Western Blot Control.
Storage	-20 C
GENE INFORMATION	
Gene Name	Cxcl10 chemokine (C-X-C motif) ligand 10 [<i>Mus musculus</i>]
Official Symbol	CXCL10
Synonyms	CXCL10; chemokine (C-X-C motif) ligand 10; C-X-C motif chemokine 10; gamma-IP10; interferon activated gene 10; small-inducible cytokine B10; interferon-gamma induced protein CRG-2; interferon-gamma-induced protein CRG-2; 10 kDa interferon gamma-induced protein; small inducible cytokine B subfamily (Cys-X-Cys), member 10; C7; IP10; CRG-2; INP10; IP-10; Ifi10; mob-1; Scyb10; gIP-10;
Gene ID	15945
mRNA Refseq	NM_021274
Protein Refseq	NP_067249
Pathway	Chemokine receptors bind chemokines, organism-specific biosystem; Chemokine

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA



signaling pathway, organism-specific biosystem; Chemokine signaling pathway, conserved biosystem; Class A/1 (Rhodopsin-like receptors), organism-specific biosystem; Cytokine-cytokine receptor interaction, organism-specific biosystem; Cytokine-cytokine receptor interaction, conserved biosystem; Cytosolic DNA-sensing pathway, organism-specific biosystem;

Function

chemokine activity; cytokine activity;

Tel: 1-631-559-9269 1-516-512-3133

Email: info@creative-biomart.com Fax: 1-631-938-8127

45-1 Ramsey Road, Shirley, NY 11967, USA