

Recombinant Human EPCAM Protein, His-tagged, Biotinylated

Cat. No. EPCAM-220H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Biotinylated Recombinant human EPCAM protein with His tag was expressed in HEK293..
Species	Human
Source	HEK293
ProteinLength	314
Description	This gene encodes a carcinoma-associated antigen and is a member of a family that includes at least two type I membrane proteins. This antigen is expressed on most normal epithelial cells and gastrointestinal carcinomas and functions as a homotypic calcium-independent cell adhesion molecule. The antigen is being used as a target for immunotherapy treatment of human carcinomas. Mutations in this gene result in congenital tufting enteropathy.
Form	Lyophilized
Molecular Mass	29 kDa
AA Sequence	MAPPQVLAFLGLLLAAATATFAAAQEEVCENYKLAVNCFVNNNRQCQCTSVGAQNT VICKSLAAKCLVMKAEMNGSKLGRRRAKPEGALQNNNDGLYDPDCDESGLFKAKQCN GTSMCWCVNNTAGVRRRTDKDTEITCSERVRTYWIIELKHKAREKPYDSKSLRTALQK EITTRYQLDPKFITSILYENNVITIDLQNSSQKTQNDVDIADVAYYFEKDKGSLFHS KKMDLTVNGEQLDLDPGQTLIYYVDEKAPEFSMQGLKAGVIAVIVVVVIAVVAGIVVL

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

	VISRKKRMAKYEKAEIKEMGEMHRELNA
Purity	> 98%
Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Concentration	1 mg/mL
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered colorless solution.
Reconstitution	Reconstitute in sterile distilled H2O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.
Conjugation	Biotin

GENE INFORMATION

Gene Name	EPCAM epithelial cell adhesion molecule [Homo sapiens (human)]
Official Symbol	EPCAM
Synonyms	EPCAM; epithelial cell adhesion molecule; antigen identified by monoclonal AUA1 , M4S1, MIC18, TACSTD1, tumor associated calcium signal transducer 1; 17 1A;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

323/A3; CD326; CO 17A; EGP 2; EGP34; EGP40; Ep CAM; ESA; GA733 2; HEA125; KS1/4; KSA; Ly74; MH99; MK 1; MOC31; TACST 1; TROP1; epithelial glycoprotein 314; human epithelial glycoprotein-2; cell surface glycoprotein Trop-1; adenocarcinoma-associated antigen; tumor-associated calcium signal transducer 1; major gastrointestinal tumor-associated protein GA733-2; membrane component, chromosome 4, surface marker (35kD glycoprotein); M4S1; MK-1; DIAR5; EGP-2; MIC18; EGP314; HNPCC8; TACSTD1; GA733-2;

Gene ID	4072
mRNA Refseq	NM_002354
Protein Refseq	NP_002345
MIM	185535
UniProt ID	P16422