

Recombinant Human FCGR2A Protein (H167)

Cat. No. FCGR2A-543H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human FCGR2A protein (H167) without tag was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	317
Description	This gene encodes one member of a family of immunoglobulin Fc receptor genes found on the surface of many immune response cells. The protein encoded by this gene is a cell surface receptor found on phagocytic cells such as macrophages and neutrophils, and is involved in the process of phagocytosis and clearing of immune complexes. Alternative splicing results in multiple transcript variants.
Form	Lyophilized
AA Sequence	MTMETQMSQNVCPRLWLLQPLTVLLLLASADSQAAAPPKAVLKLEPPWINVLQED SVTLTCQGARSPESDSIQWFHNGNLIPTHTQPSYRFKANNNDSGEYTCQTGQTSLS DPVHLTVLSEWLVLQTPHLEFQEGETIMLRCHSWKDKPLVKVTFQNGKSQKFSLH DPTFSIPQANHSHSGDYHCTGNIGYTLFSSKPVTITVQVPSMGSSSPMGIIVAVVIATA VAAIVA AVVALIYCRKKRISANSTDPVKAAQFEPGRQMI AIRKRQLEETNNDYETAD GGYMTLNPRAPTDDDKNIYLTLPNDHVNNSNN
Purity	> 98%

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Concentration	1 mg/mL
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered colorless solution.
Reconstitution	Reconstitute in sterile distilled H2O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.

GENE INFORMATION

Gene Name	FCGR2A Fc fragment of IgG, low affinity IIa, receptor (CD32) [Homo sapiens (human)]
Official Symbol	FCGR2A
Synonyms	FCGR2A; Fc fragment of IgG, low affinity IIa, receptor (CD32); Fc fragment of IgG, low affinity IIa, receptor for (CD32) , FCG2, FCGR2, FCGR2A1; low affinity immunoglobulin gamma Fc region receptor II-a; CD32; CD32A; CDw32; IGFR2; Immunoglobulin G Fc receptor II; fcRII-a; fc-gamma-RIIa; fc-gamma RII-a; igG Fc receptor II-a; FCG2; FcGR; FCGR2; FCGR2A1; MGC23887; MGC30032;
Gene ID	2212

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_001136219
Protein Refseq	NP_001129691
MIM	146790
UniProt ID	P12318

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA