

Recombinant Human CD70 Protein, Fc-tagged

Cat. No. CD70-623H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human CD70 protein with Fc tag was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	193
Description	The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis.
Form	Lyophilized
Molecular Mass	44.6 kDa
AA Sequence	MPEEGSGCSVRRRPYGCVLRAALVPLVAGLVICLVVCIQRFAQAQQQLPLESLGWD VAELQLNHTGPQQDPRLYWQGGPALGRSFLHGPGLDKGQLRIHRDGIYMVHIQVTL AICSSTTASRHPTTLAVGICSPASRSISLLRLSFHQGCTIASQRLTPLARGDTLCTNL TGTLLPSRNTDETFFGVQWVRP

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	> 98%
Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Concentration	1 mg/mL
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered colorless solution.
Reconstitution	Reconstitute in sterile distilled H2O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.

GENE INFORMATION

Gene Name	CD70 CD70 molecule [Homo sapiens (human)]
Official Symbol	CD70
Synonyms	CD70; CD70 molecule; CD27LG, TNFSF7, tumor necrosis factor (ligand) superfamily, member 7; CD70 antigen; CD27L; CD27-L; CD27 ligand; Ki-24 antigen; surface antigen CD70; tumor necrosis factor ligand superfamily member 7; tumor necrosis factor (ligand) superfamily, member 7; CD27LG; TNFSF7;
Gene ID	970

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

mRNA Refseq	NM_001252
Protein Refseq	NP_001243
MIM	602840
UniProt ID	P32970

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA