

Recombinant Cynomolgus monkey CD40 Protein, Fc-tagged

Cat. No. CD40-627C Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant Cynomolgus monkey CD40 protein with Fc tag was expressed in HEK293.
Species	Cynomolgus
Source	HEK293
ProteinLength	278
Description	This gene is a member of the TNF-receptor superfamily. The encoded protein is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. Mutations affecting this gene are the cause of autosomal recessive hyper-IgM immunodeficiency type 3 (HIGM3). Multiple alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.
Form	Lyophilized

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Molecular Mass	45.8 kDa
AA Sequence	MVRLPLQCVLWGCLLTAVYPEPPTACREKQYLINSQCCSLCQPGQKLVS DCTEFTE TECLPCGESEFLDTWNRETRCHQHKYCDPNLGLRVQQKGTSETDTICTCEEGLHCT SESCESCVPHRSCLPFGVGKQIATGVSDTICEPCPVGFFSNVSSAFEKCRPWTSCE TKDLVVQQAGTNKTDVVCGPQDRQRALVVIPICLGILFVILLVLVFIKKVAKKPNDKV PHPKQEPQEINFPDDLPGSNPAAPVQETLHGCQPVTQEDGKESRISVQERQ
Purity	> 98%
Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Concentration	1 mg/mL
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered colorless solution.
Reconstitution	Reconstitute in sterile distilled H ₂ O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.

GENE INFORMATION

Gene Name	CD40 CD40 molecule, TNF receptor superfamily member 5 [Macaca mulatta (Rhesus monkey)]
------------------	--

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Official Symbol	CD40
Synonyms	CD40; CD40 molecule, TNF receptor superfamily member 5; tumor necrosis factor receptor superfamily member 5;
Gene ID	707749
mRNA Refseq	NM_001265862
Protein Refseq	NP_001252791
UniProt ID	F6SNI1

 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA