

Recombinant Human TNFRSF18 Protein

Cat. No. TNFRSF18-661H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human TNFRSF18 protein without tag was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	241
Description	This gene encodes a member of the TNF-receptor superfamily. The encoded receptor has been shown to have increased expression upon T-cell activation, and it is thought to play a key role in dominant immunological self-tolerance maintained by CD25(+)CD4(+) regulatory T cells. Knockout studies in mice also suggest the role of this receptor is in the regulation of CD3-driven T-cell activation and programmed cell death. Three alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.
Form	Lyophilized
Molecular Mass	15.4 kDa
AA Sequence	MAQHGMGAFRALCGLALLCALSLGQRPTGGPGCGPGRLLLGTGTDARCCRVHTT RCCRDYPGEECCSEWDCMCVQPEFHCGDPCCTTCRHHPCPPGQGVQSQGKFSF GFQCIDCASGTFSGGHEGHCKPWTDCQFGFLTVPNGKTHNAVCPGSPPAEPL GWLTVVLLAVAACVLLLTSAQLGLHIWQLRSQCMWPRETQLLLEVPPSTEDARSCQ FPEEERGERSAEEKGRLGDLWV

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	> 98%
Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Concentration	1 mg/mL
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered colorless solution.
Reconstitution	Reconstitute in sterile distilled H2O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.

GENE INFORMATION

Gene Name	TNFRSF18 tumor necrosis factor receptor superfamily, member 18 [Homo sapiens (human)]
Official Symbol	TNFRSF18
Synonyms	TNFRSF18; tumor necrosis factor receptor superfamily, member 18; tumor necrosis factor receptor superfamily member 18; AITR; CD357; GITR; activation-inducible TNFR family receptor; glucocorticoid-induced TNFR-related protein; TNF receptor superfamily activation-inducible protein; GITR-D;

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Gene ID	8784
mRNA Refseq	NM_004195
Protein Refseq	NP_004186
MIM	603905
UniProt ID	Q9Y5U5

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA