

Recombinant Human TNFRSF4 Protein

Cat. No. TNFRSF4-680H **Lot. No.** (See product label)

SPECIFICATION

Product Overview	Recombinant human TNFRSF4 protein without tag was expressed in HEK293.
Species	Human
Source	HEK293
ProteinLength	277
Description	The protein encoded by this gene is a member of the TNF-receptor superfamily. This receptor has been shown to activate NF-kappaB through its interaction with adaptor proteins TRAF2 and TRAF5. Knockout studies in mice suggested that this receptor promotes the expression of apoptosis inhibitors BCL2 and BCL2L1/BCL2-XL, and thus suppresses apoptosis. The knockout studies also suggested the roles of this receptor in CD4+ T cell response, as well as in T cell-dependent B cell proliferation and differentiation.
Form	Lyophilized
Molecular Mass	22 kDa
AA Sequence	MCVGARRLGRGPCAALLLLGLGLSTVTGLHCVGDTYPSNDRCCHECRPGNGMVSR CSRSQNTVCRPCGPGFYNDVVSSKPKPCTWCNLRSGSERKQLCTATQDTVCR RAGTQPLDSYKPGVDCAPCPPGHFSPGDNQACKPWTNCTLAGKHTLQPASNSSDA ICEDRDPPATQPQETQGPPARPITVQPTAWPRTSQQGPSTRPVEVPGGRAVAAILG LGLVLGLLGPLAILLALYLLRRDQRLPPDAHKPPGGGSFRTPIQEEQADAHSTLAKI

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

Purity	> 98%
Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Concentration	1 mg/mL
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered colorless solution.
Reconstitution	Reconstitute in sterile distilled H2O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.

GENE INFORMATION

Gene Name	TNFRSF4 tumor necrosis factor receptor superfamily, member 4 [Homo sapiens (human)]
Official Symbol	TNFRSF4
Synonyms	TNFRSF4; tumor necrosis factor receptor superfamily, member 4; TXGP1L; tumor necrosis factor receptor superfamily member 4; ACT35; CD134; OX40; OX40 antigen; ACT35 antigen; ATC35 antigen; CD134 antigen; OX40 homologue; OX40L receptor; OX40 cell surface antigen; lymphoid activation antigene ACT35; TAX transcriptionally-activated glycoprotein 1 receptor; tax-transcriptionally activated

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

glycoprotein 1 receptor;

Gene ID 7293**mRNA Refseq** NM_003327**Protein Refseq** NP_003318**MIM** 600315**UniProt ID** P43489 Tel: 1-631-559-9269 1-516-512-3133 Email: info@creative-biomart.com  Fax: 1-631-938-8127 45-1 Ramsey Road, Shirley, NY 11967, USA