

Recombinant Human SIRPA Protein, Fc-tagged

Cat. No. SIRPA-723H Lot. No. (See product label)

SPECIFICATION

Product Overview	Recombinant human SIRPA protein with Fc tag was expressed in CHO
Species	Human
Source	CHO
ProteinLength	504
Description	The protein encoded by this gene is a member of the signal-regulatory-protein (SIRP) family, and also belongs to the immunoglobulin superfamily. SIRP family members are receptor-type transmembrane glycoproteins known to be involved in the negative regulation of receptor tyrosine kinase-coupled signaling processes. This protein can be phosphorylated by tyrosine kinases. The phospho-tyrosine residues of this PTP have been shown to recruit SH2 domain containing tyrosine phosphatases (PTP), and serve as substrates of PTPs. This protein was found to participate in signal transduction mediated by various growth factor receptors. CD47 has been demonstrated to be a ligand for this receptor protein. This gene and its product share very high similarity with several other members of the SIRP family. These related genes are located in close proximity to each other on chromosome 20p13. Multiple alternatively spliced transcript variants have been determined for this gene.
Form	Lyophilized
Molecular Mass	63.5 kDa

 Tel: 1-631-559-9269 1-516-512-3133

 Email: info@creative-biomart.com  Fax: 1-631-938-8127

 45-1 Ramsey Road, Shirley, NY 11967, USA

AA Sequence	MEPAGPAPGRLGPLLCLLLAASCAWSGVAGEEELQVIQPDKSVLVAAGETATLRCT ATSLIPVGPIQWFRGAGPGRELIYNQKEGHFPRVTTVSDLTKRNNMDFSIRIGNITPA DAGTYCYVKFRKGGSPDDVEFKSGAGTELSVRAKPSAPVVSGPAARATPQHTVSFTC ESHGFSRPDITLKWFKNGNELSDFQTNVDPVGESVSYSIHSTAKVVLTREDVHSQVI CEVAHVTLQGDPLRGTANLSETIRVPPTLEVTQQPVRAENQVNVTCQVRKFYPQRL QLTWLENGNVSRTEASTVTENKDGTYNWMSWLLVNVSAHRDDVKLTCQVEHDG QPAVSKSHDLKVS AHPKEQGSNTAAENTGSNERNIYIVGVVCTLLVALLMAALYLV RIRQKKAQGSTSSTRLHEPEKNAREITQDTNDITYADLNLPKGKKPAPQAAEPNNHT EYASIQTSPQPASEDTLTADLDMVHLNRTPKQPAPKPEPSFSEYASVQVPRK
Purity	> 98%
Applications	WB; ELISA; FACS; FC
Stability	This bioreagent is stable at 4 centigrade (short-term) and -70 centigrade(long-term). After reconstitution, sample may be stored at 4 centigrade for 2-7 days and below -18 centigrade for future use.
Storage	At -20 centigrade.
Concentration	1 mg/mL
Storage Buffer	PBS (pH 7.4-7.5). Sterile-filtered colorless solution.
Reconstitution	Reconstitute in sterile distilled H₂O to no less than 100 µg/mL; dilute reconstituted stock further in other aqueous solutions if needed. Please review COA for lot-specific instructions. Final measurements should be determined by the end-user for optimal performance.

GENE INFORMATION

Gene Name	SIRPA signal-regulatory protein alpha [Homo sapiens (human)]
Official Symbol	SIRPA
Synonyms	SIRPA; signal-regulatory protein alpha; protein tyrosine phosphatase, non receptor type substrate 1 , PTPNS1; tyrosine-protein phosphatase non-receptor type substrate 1; BIT; CD172a; MFR; MYD 1; P84; SHPS 1; SHPS1; SIRP; SIRP ALPHA 1; SIRPalpha; SIRPalpha2; myd-1 antigen; inhibitory receptor SHPS-1; macrophage fusion receptor; CD172 antigen-like family member A; tyrosine phosphatase SHP substrate 1; brain-immunoglobulin-like molecule with tyrosine-based activation motifs; MYD-1; CD172A; PTPNS1;
Gene ID	140885
mRNA Refseq	NM_001040022
Protein Refseq	NP_001035111
MIM	602461
UniProt ID	P78324